

1. Schafer, W., et al. Good urodynamic practices: uroflowmetry, filling cystometry, and pressure-flow studies. *Neurourol Urodyn*, 2002. 21: 261. <http://www.ncbi.nlm.nih.gov/pubmed/11948720>
2. Abrams, P., et al. Reviewing the ICS 2002 terminology report: the ongoing debate. *Neurourol, Urodyn*, 2009. 28: 287. <http://www.ncbi.nlm.nih.gov/pubmed/19350662>
3. Abrams, P., et al. The standardisation of terminology of lower urinary tract function: report from the Standardisation Subcommittee of the International Continence Society. *Neurourol Urodyn*, 2002. 21: 167. <http://www.ncbi.nlm.nih.gov/pubmed/11857671>
4. Groen, J., et al. Summary of European Association of Urology (EAU) Guidelines on Neuro-Urology. *Eur Urol*, 2015. <http://www.ncbi.nlm.nih.gov/pubmed/26194043>
5. Schneider, M.P., et al. Tibial Nerve Stimulation for Treating Neurogenic Lower Urinary Tract Dysfunction: A Systematic Review. *Eur Urol*, 2015. 68(5): p. 859-67 <http://www.ncbi.nlm.nih.gov/pubmed/26194043>
6. Nosseir, M., et al. Clinical usefulness of urodynamic assessment for maintenance of bladder function in patients with spinal cord injury. *Neurourol Urodyn*, 2007. 26: 228. <http://www.ncbi.nlm.nih.gov/pubmed/16998859>
7. Panicker, J.N., et al. Lower urinary tract dysfunction in the neurological patient: clinical assessment and management. *Lancet Neurol*, 2015. 14:720. <http://www.ncbi.nlm.nih.gov/pubmed/26067125>
8. Kessler, T.M., et al. Transcutaneous electrical nerve stimulation and percutaneous tibial nerve stimulation for neurogenic lower urinary tract dysfunction: a systematic review. *PROSPERO*, 2014. [http://www.crd.york.ac.uk/PROSPERO/display\\_record.asp?ID=CRD42014008678](http://www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42014008678)
9. 't Hoen, L., et al. Which measures are available to evaluate sexual function/dysfunction in adult neuro-urological patients and which are the most appropriate? A systematic review. *PROSPERO*, 2014. [http://www.crd.york.ac.uk/PROSPERO/display\\_record.asp?ID=CRD42014015287](http://www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42014015287)
10. Phillips B, et al. Oxford Centre for Evidence-based Medicine Levels of Evidence. Updated by Jeremy Howick March 2009. 1998. <http://www.cebm.net/oxford-centre-evidence-based-medicine-levels-evidence-march-2009/>
11. Townsend, N., et al. Cardiovascular disease in Europe - epidemiological update 2015. *Eur Heart, J*, 2015. <http://www.ncbi.nlm.nih.gov/pubmed/26306399>
12. Tibaek, S., et al. Prevalence of lower urinary tract symptoms (LUTS) in stroke patients: a cross-sectional, clinical survey. *Neurourol Urodyn*, 2008. 27: 763. <http://www.ncbi.nlm.nih.gov/pubmed/18551565>
13. Marinkovic, S.P., et al. Voiding and sexual dysfunction after cerebrovascular accidents. *J Urol*, 2001. 165: 359. <http://www.ncbi.nlm.nih.gov/pubmed/11176374>
14. Rotar, M., et al. Stroke patients who regain urinary continence in the first week after acute first-ever stroke have better prognosis than patients with persistent lower urinary tract dysfunction. *Neurourol Urodyn*, 2011. 30: 1315. <http://www.ncbi.nlm.nih.gov/pubmed/21488096>
15. Lobo, A., et al. Prevalence of dementia and major subtypes in Europe: A collaborative study of population-based cohorts. *Neurologic Diseases in the Elderly Research Group. Neurology*, 2000. 54:S4. <http://www.ncbi.nlm.nih.gov/pubmed/10854354>
16. Na, H.R., et al. Urinary incontinence in Alzheimer's disease is associated with Clinical Dementia Rating-Sum of Boxes and Barthel Activities of Daily Living. *Asia Pac Psychiatry*, 2015. 7: 113. <http://www.ncbi.nlm.nih.gov/pubmed/23857871>
17. Grant, R.L., et al. First diagnosis and management of incontinence in older people with and without dementia in primary care: a cohort study using The Health Improvement Network primary care database. *PLoS Med*, 2013. 10: e1001505. <http://www.ncbi.nlm.nih.gov/pubmed/24015113>
18. Pringsheim, T., et al. The prevalence of Parkinson's disease: a systematic review and meta-analysis. *Mov Disord*, 2014. 29: 1583. <http://www.ncbi.nlm.nih.gov/pubmed/24976103>
19. Ragab, M.M., et al. Idiopathic Parkinson's disease patients at the urologic clinic. *Neurourol Urodyn*, 2011. 30: 1258. <http://www.ncbi.nlm.nih.gov/pubmed/21404318>
20. Papatsoris, A.G., et al. Urinary and erectile dysfunction in multiple system atrophy (MSA). *Neurourol Urodyn*, 2008. 27: 22. <http://www.ncbi.nlm.nih.gov/pubmed/17563111>
21. Kim, M., et al. Impaired detrusor contractility is the pathognomonic urodynamic finding of multiple system atrophy compared to idiopathic Parkinson's disease. *Parkinsonism Relat Disord*, 2015. 21:205. <http://www.ncbi.nlm.nih.gov/pubmed/25534084>
22. Sakakibara, R., et al. A guideline for the management of bladder dysfunction in Parkinson's disease and other gait disorders. *Neurourol Urodyn*, 2015. <http://www.ncbi.nlm.nih.gov/pubmed/25810035>
23. Dolecek, T.A., et al. CBTRUS statistical report: primary brain and central nervous system tumors diagnosed in the United States in 2005-2009. *Neuro Oncol*, 2012. 14 Suppl 5: v1. <http://www.ncbi.nlm.nih.gov/pubmed/23095881>
24. Maurice-Williams, R.S. Micturition symptoms in frontal tumours. *J Neurol Neurosurg Psychiatry*, 1974. 37: 431. <http://www.ncbi.nlm.nih.gov/pubmed/4365244>
25. Christensen, D., et al. Prevalence of cerebral palsy, co-occurring autism spectrum disorders, and motor functioning - Autism and Developmental Disabilities Monitoring Network, USA, 2008. *Dev Med Child Neurol*, 2014. 56: 59. <http://www.ncbi.nlm.nih.gov/pubmed/24117446>
26. Marciniak, C., et al. Urinary incontinence in adults with cerebral palsy: prevalence, type, and effects on participation. *PM R*, 2014. 6: 110. <http://www.ncbi.nlm.nih.gov/pubmed/23978464>



27. Singh, A., et al. Global prevalence and incidence of traumatic spinal cord injury. *Clin Epidemiol*, 2014. 6: 309. <http://www.ncbi.nlm.nih.gov/pubmed/25278785>
28. Weld, K.J., et al. Association of level of injury and bladder behavior in patients with post-traumatic spinal cord injury. *Urology*, 2000. 55: 490. <http://www.ncbi.nlm.nih.gov/pubmed/10736489>
29. Kondo, A., et al. Neural tube defects: prevalence, etiology and prevention. *Int J Urol*, 2009. 16: 49. <http://www.ncbi.nlm.nih.gov/pubmed/19120526>
30. Sawin, K.J., et al. The National Spina Bifida Patient Registry: profile of a large cohort of participants from the first 10 clinics. *J Pediatr*, 2015. 166: 444. <http://www.ncbi.nlm.nih.gov/pubmed/25444012>
31. Bartolin, Z., et al. Relationship between clinical data and urodynamic findings in patients with lumbar intervertebral disk protrusion. *Urol Res*, 2002. 30: 219. <http://www.ncbi.nlm.nih.gov/pubmed/12202938>
32. Lange, M.M., et al. Urinary and sexual dysfunction after rectal cancer treatment. *Nat Rev Urol*, 2011. 8: 51. <http://www.ncbi.nlm.nih.gov/pubmed/21135876>
33. Federation, I.D., *IDF Diabetes Atlas, 6th edn.* 2013, International Diabetes Federation: Brussels, Belgium.
34. Yuan, Z., et al. Diabetic cystopathy: A review. *J Diabetes*, 2015. 7: 442. <http://www.ncbi.nlm.nih.gov/pubmed/25619174>
35. Pugliatti, M., et al. The epidemiology of multiple sclerosis in Europe. *Eur J Neurol*, 2006. 13: 700. <http://www.ncbi.nlm.nih.gov/pubmed/16834700>
36. de Seze, M., et al. The neurogenic bladder in multiple sclerosis: review of the literature and proposal of management guidelines. *Mult Scler*, 2007. 13: 915. <http://www.ncbi.nlm.nih.gov/pubmed/17881401>
37. Krassioukov, A., et al. International standards to document remaining autonomic function after spinal cord injury. *J Spinal Cord Med*, 2012. 35: 201. <http://www.ncbi.nlm.nih.gov/pubmed/229257462>
38. Irwin, D.E., et al. Understanding the elements of overactive bladder: questions raised by the EPIC study. *BJU Int*, 2008. 101: 1381. <http://www.ncbi.nlm.nih.gov/pubmed/18336602>
39. Fix, J.D., *Neuroanatomy.* 4th ed. 2008, Philadelphia, Pennsylvania, USA.
40. Maynard, F.M., Jr., et al. International Standards for Neurological and Functional Classification of Spinal Cord Injury. American Spinal Injury Association. *Spinal Cord*, 1997. 35: 266. <http://www.ncbi.nlm.nih.gov/pubmed/9160449>
41. Klevmark, B. Natural pressure-volume curves and conventional cystometry. *Scand J Urol Nephrol Suppl*, 1999. 201: 1. <http://www.ncbi.nlm.nih.gov/pubmed/10573769>
42. Del Popolo, G., et al. Diagnosis and therapy for neurogenic bladder dysfunctions in multiple sclerosis patients. *Neurol Sci*, 2008. 29 Suppl 4: S352. <http://www.ncbi.nlm.nih.gov/pubmed/19089675>
43. Satar, N., et al. The effects of delayed diagnosis and treatment in patients with an occult spinal dysraphism. *J Urol*, 1995. 154: 754. <http://www.ncbi.nlm.nih.gov/pubmed/7609171>
44. Watanabe, T., et al. High incidence of occult neurogenic bladder dysfunction in neurologically intact patients with thoracolumbar spinal injuries. *J Urol*, 1998. 159: 965. <http://www.ncbi.nlm.nih.gov/pubmed/9474194>
45. Ahlberg, J., et al. Neurological signs are common in patients with urodynamically verified "idiopathic" bladder overactivity. *Neurourol Urodyn*, 2002. 21: 65. <http://www.ncbi.nlm.nih.gov/pubmed/11835426>
46. Bemelmans, B.L., et al. Evidence for early lower urinary tract dysfunction in clinically silent multiple sclerosis. *J Urol*, 1991. 145: 1219. <http://www.ncbi.nlm.nih.gov/pubmed/2033697>
47. Klausner, A.P., et al. The neurogenic bladder: an update with management strategies for primary care physicians. *Med Clin North Am*, 2011. 95: 111. <http://www.ncbi.nlm.nih.gov/pubmed/21095415>
48. Bors, E., et al. History and physical examination in neurological urology. *J Urol*, 1960. 83: 759. <http://www.ncbi.nlm.nih.gov/pubmed/13802958>
49. Cameron, A.P., et al. The Severity of Bowel Dysfunction in Patients with Neurogenic Bladder. *J Urol*, 2015. <http://www.ncbi.nlm.nih.gov/pubmed/25956470>
50. Vodušek, D.B. Lower urinary tract and sexual dysfunction in neurological patients. *Eur Neurol*, 2014. 72: 109. <http://www.ncbi.nlm.nih.gov/pubmed/24993182>
51. Linsmeyer, T.A., et al. Accuracy of individuals with spinal cord injury at predicting urinary tract infections based on their symptoms. *J Spinal Cord Med*, 2003. 26: 352. <http://www.ncbi.nlm.nih.gov/pubmed/14992336>
52. Massa, L.M., et al. Validity, accuracy, and predictive value of urinary tract infection signs and symptoms in individuals with spinal cord injury on intermittent catheterization. *J Spinal Cord Med*, 2009. 32: 568. <http://www.ncbi.nlm.nih.gov/pubmed/20025153>
53. Honjo, H., et al. Impact of convenience void in a bladder diary with urinary perception grade to assess overactive bladder symptoms: a community-based study. *Neurourol Urodyn*, 2010. 29: 1286. <http://www.ncbi.nlm.nih.gov/pubmed/20878998>
54. Naoemova, I., et al. Reliability of the 24-h sensation-related bladder diary in women with urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct*, 2008. 19: 955. <http://www.ncbi.nlm.nih.gov/pubmed/18235981>
55. Henze, T. Managing specific symptoms in people with multiple sclerosis. *Int MS J*, 2005. 12: 60. <http://www.ncbi.nlm.nih.gov/pubmed/16417816>
56. Liu, C.W., et al. The relationship between bladder management and health-related quality of life in patients with spinal cord injury in the UK. *Spinal Cord*, 2010. 48: 319. <http://www.ncbi.nlm.nih.gov/pubmed/19841636>
57. Pannek, J., et al. Does optimizing bladder management equal optimizing quality of life? Correlation between health-related quality of life and urodynamic parameters in patients with spinal cord lesions. *Urology*, 2009. 74: 263. <http://www.ncbi.nlm.nih.gov/pubmed/19428089>



58. Patel, D.P., et al. Patient reported outcomes measures in neurogenic bladder and bowel: A systematic review of the current literature. *Neurourol Urodyn*, 2014. <http://www.ncbi.nlm.nih.gov/pubmed/25327455>
59. Bonniaud, V., et al. Qualiveen, a urinary-disorder specific instrument: 0.5 corresponds to the minimal important difference. *J Clin Epidemiol*, 2008. 61: 505. <http://www.ncbi.nlm.nih.gov/pubmed/18394545>
60. Bonniaud, V., et al. Development and validation of the short form of a urinary quality of life questionnaire: SF-Qualiveen. *J Urol*, 2008. 180: 2592. <http://www.ncbi.nlm.nih.gov/pubmed/18950816>
61. Bonniaud, V., et al. Italian version of Qualiveen-30: cultural adaptation of a neurogenic urinary disorder-specific instrument. *Neurourol Urodyn*, 2011. 30: 354. <http://www.ncbi.nlm.nih.gov/pubmed/21305589>
62. Ciudin, A., et al. Quality of life of multiple sclerosis patients: translation and validation of the Spanish version of Qualiveen. *Neurourol Urodyn*, 2012. 31: 517. <http://www.ncbi.nlm.nih.gov/pubmed/22396437>
63. D'Ancona, C.A., et al. Quality of life of neurogenic patients: translation and validation of the Portuguese version of Qualiveen. *Int Urol Nephrol*, 2009. 41: 29. <http://www.ncbi.nlm.nih.gov/pubmed/18528780>
64. Pannek, J., et al. [Quality of life in German-speaking patients with spinal cord injuries and bladder dysfunctions. Validation of the German version of the Qualiveen questionnaire]. *Urologe A*, 2007. 46: 1416. <http://www.ncbi.nlm.nih.gov/pubmed/17605119>
65. Welk, B., et al. The conceptualization and development of a patient-reported neurogenic bladder symptom score. *Res Rep Urol*, 2013. 5: 129. <http://www.ncbi.nlm.nih.gov/pubmed/24400244>
66. Gulick, E.E. Bowel management related quality of life in people with multiple sclerosis: psychometric evaluation of the QoL-BM measure. *Int J Nurs Stud*, 2011. 48: 1066. <http://www.ncbi.nlm.nih.gov/pubmed/21377677>
67. Tsang, B., et al. A systematic review and comparison of questionnaires in the management of spinal cord injury, multiple sclerosis and the neurogenic bladder. *Neurourol Urodyn*, 2015. <http://www.ncbi.nlm.nih.gov/pubmed/25620137>
68. Schurch, B., et al. Reliability and validity of the Incontinence Quality of Life questionnaire in patients with neurogenic urinary incontinence. *Arch Phys Med Rehabil*, 2007. 88: 646. <http://www.ncbi.nlm.nih.gov/pubmed/17466735>
69. Hollingworth, W., et al. Exploring the impact of changes in neurogenic urinary incontinence frequency and condition-specific quality of life on preference-based outcomes. *Qual Life Res*, 2010. 19: 323. <http://www.ncbi.nlm.nih.gov/pubmed/20094804>
70. Cella, D.F., et al. Validation of the functional assessment of multiple sclerosis quality of life instrument. *Neurology*, 1996. 47: 129. <http://www.ncbi.nlm.nih.gov/pubmed/8710066>
71. Wesson, J.M., et al. The functional index for living with multiple sclerosis: development and validation of a new quality of life questionnaire. *Mult Scler*, 2009. 15: 1239. <http://www.ncbi.nlm.nih.gov/pubmed/19737850>
72. Gold, S.M., et al. Disease specific quality of life instruments in multiple sclerosis: validation of the Hamburg Quality of Life Questionnaire in Multiple Sclerosis (HAQUAMS). *Mult Scler*, 2001. 7: 119. <http://www.ncbi.nlm.nih.gov/pubmed/11424632>
73. Goodin, D.S. A questionnaire to assess neurological impairment in multiple sclerosis. *Mult Scler*, 1998. 4: 444. <http://www.ncbi.nlm.nih.gov/pubmed/9839306>
74. Foley, F.W., et al. The Multiple Sclerosis Intimacy and Sexuality Questionnaire - re-validation and development of a 15-item version with a large US sample. *Mult Scler*, 2013. 19: 1197. <http://www.ncbi.nlm.nih.gov/pubmed/23369892>
75. Sanders, A.S., et al. The Multiple Sclerosis Intimacy and Sexuality Questionnaire-19 (MSISQ-19). *Sexuality and Disability*, 2000. 18: 3.
76. Marrie, R.A., et al. Validity and reliability of the MSQLI in cognitively impaired patients with multiple sclerosis. *Mult Scler*, 2003. 9: 621. <http://www.ncbi.nlm.nih.gov/pubmed/14664477>
77. Vickrey, B.G., et al. A health-related quality of life measure for multiple sclerosis. *Qual Life Res*, 1995. 4: 187. <http://www.ncbi.nlm.nih.gov/pubmed/7613530>
78. Honan, C.A., et al. The multiple sclerosis work difficulties questionnaire (MSWDQ): development of a shortened scale. *Disabil Rehabil*, 2014. 36: 635. <http://www.ncbi.nlm.nih.gov/pubmed/23786346>
79. Welk, B., et al. The validity and reliability of the neurogenic bladder symptom score. *J Urol*, 2014. 192: 452. <http://www.ncbi.nlm.nih.gov/pubmed/24518764>
80. Bonniaud, V., et al. Measuring quality of life in multiple sclerosis patients with urinary disorders using the Qualiveen questionnaire. *Arch Phys Med Rehabil*, 2004. 85: 1317. <http://www.ncbi.nlm.nih.gov/pubmed/15295759>
81. Franceschini, M., et al. Follow-up in persons with traumatic spinal cord injury: questionnaire reliability. *Eura Medicophys*, 2006. 42: 211. <http://www.ncbi.nlm.nih.gov/pubmed/17039217>
82. Noreau, L., et al. Development and assessment of a community follow-up questionnaire for the Rick Hansen spinal cord injury registry. *Arch Phys Med Rehabil*, 2013. 94: 1753. <http://www.ncbi.nlm.nih.gov/pubmed/23529142>
83. Liu, N., et al. Iatrogenic urological triggers of autonomic dysreflexia: a systematic review. *Spinal Cord*, 2015. 53: 500. <http://www.ncbi.nlm.nih.gov/pubmed/25800696>
84. Labat, J.J., et al. Diagnostic criteria for pudendal neuralgia by pudendal nerve entrapment (Nantes criteria). *Neurourol Urodyn*, 2008. 27: 306. <http://www.ncbi.nlm.nih.gov/pubmed/17828787>
85. Brown, D.L., Atlas of regional anesthesia. 3rd. ed. 2006, Philadelphia
86. Gray's anatomy, . 40th ed, ed. S. S. 2008.



87. Bellucci, C.H., et al. Neurogenic lower urinary tract dysfunction—do we need same session repeat urodynamic investigations? *J Urol*, 2012. 187: 1318. <http://www.ncbi.nlm.nih.gov/pubmed/22341264>
88. Walter, M., et al. Autonomic dysreflexia and repeatability of cardiovascular changes during same session repeat urodynamic investigation in women with spinal cord injury. *World J Urol*, 2015. <http://www.ncbi.nlm.nih.gov/pubmed/26055644>
89. Gammie, A., et al. International Continence Society guidelines on urodynamic equipment performance. *Neurourol Urodyn*, 2014. 33: 370. <http://www.ncbi.nlm.nih.gov/pubmed/24390971>
90. McGuire, E.J., et al. Leak-point pressures. *Urol Clin North Am*, 1996. 23: 253. <http://www.ncbi.nlm.nih.gov/pubmed/8659025>
91. Ozkan, B., et al. Which factors predict upper urinary tract deterioration in overactive neurogenic bladder dysfunction? *Urology*, 2005. 66: 99. <http://www.ncbi.nlm.nih.gov/pubmed/15992868>
92. Wang, Q.W., et al. Is it possible to use urodynamic variables to predict upper urinary tract dilatation in children with neurogenic bladder-sphincter dysfunction? *BJU Int*, 2006. 98: 1295. <http://www.ncbi.nlm.nih.gov/pubmed/17034510>
93. Linsenmeyer, T.A., et al. The impact of urodynamic parameters on the upper tracts of spinal cord injured men who void reflexly. *J Spinal Cord Med*, 1998. 21: 15. <http://www.ncbi.nlm.nih.gov/pubmed/9541882>
94. McGuire, E.J., et al. Prognostic value of urodynamic testing in myelodysplastic patients. *J Urol*, 1981. 126: 205. <http://www.ncbi.nlm.nih.gov/pubmed/7196460>
95. Krongrad, A., et al. Bladder neck dysynergia in spinal cord injury. *Am J Phys Med Rehabil*, 1996. 75:204. <http://www.ncbi.nlm.nih.gov/pubmed/8663928>
96. Weld, K.J., et al. Clinical significance of detrusor sphincter dyssynergia type in patients with posttraumatic spinal cord injury. *Urology*, 2000. 56: 565. <http://www.ncbi.nlm.nih.gov/pubmed/11018603>
97. Rossier, A.B., et al. 5-microtransducer catheter in evaluation of neurogenic bladder function. *Urology*, 1986. 27: 371. <http://www.ncbi.nlm.nih.gov/pubmed/3962062>
98. Al-Ali, M., et al. A 10 year review of the endoscopic treatment of 125 spinal cord injured patients with vesical outlet obstruction: does bladder neck dyssynergia exist? *Paraplegia*, 1996. 34: 34. <http://www.ncbi.nlm.nih.gov/pubmed/8848321>
99. Bacsu, C.D., et al. Diagnosing detrusor sphincter dyssynergia in the neurological patient. *BJU Int*, 2012. 109 Suppl 3: 31. <http://www.ncbi.nlm.nih.gov/pubmed/22458490>
100. Lose, G., et al. Standardisation of urethral pressure measurement: report from the Standardisation Sub-Committee of the International Continence Society. *Neurourol Urodyn*, 2002. 21: 258. <http://www.ncbi.nlm.nih.gov/pubmed/11948719>
101. Marks, B.K., et al. Videourodynamics: indications and technique. *Urol Clin North Am*, 2014. 41: 383. <http://www.ncbi.nlm.nih.gov/pubmed/25063594>
102. Geirsson, G., et al. The ice-water test—a simple and valuable supplement to routine cystometry. *Br J Urol*, <http://www.ncbi.nlm.nih.gov/pubmed/8343894> 1993. 71: 681
103. Geirsson, G., et al. Pressure, volume and infusion speed criteria for the ice-water test. *Br J Urol*, 1994. 73: 498. <http://www.ncbi.nlm.nih.gov/pubmed/8012770>
104. Al-Hayek, S., et al. The 50-year history of the ice water test in urology. *J Urol*, 2010. 183: 1686. <http://www.ncbi.nlm.nih.gov/pubmed/20299050>
105. Lapedes, J. Neurogenic bladder. Principles of treatment. *Urol Clin North Am*, 1974. 1: 81. <http://www.ncbi.nlm.nih.gov/pubmed/4428540>
106. Riedl, C.R., et al. Electromotive administration of intravesical bethanechol and the clinical impact on acontractile detrusor management: introduction of a new test. *J Urol*, 2000. 164: 2108. <http://www.ncbi.nlm.nih.gov/pubmed/11061937>
107. Podnar, S., et al. Lower urinary tract dysfunction in patients with peripheral nervous system lesions. *Handb Clin Neurol*, 2015. 130: 203. <http://www.ncbi.nlm.nih.gov/pubmed/26003246>
108. Ouyang, L., et al. Characteristics and survival of patients with end stage renal disease and spina bifida in the United States renal data system. *J Urol*, 2015. 193: 558. <http://www.ncbi.nlm.nih.gov/pubmed/25167993>
109. Lawrenson, R., et al. Renal failure in patients with neurogenic lower urinary tract dysfunction. *Neuroepidemiology*, 2001. 20: 138. <http://www.ncbi.nlm.nih.gov/pubmed/11359083>
110. Averbek, M.A., et al. Follow-up of the neuro-urological patient: a systematic review. *BJU Int*, 2015. 115 Suppl 6: 39. <http://www.ncbi.nlm.nih.gov/pubmed/25891319>
111. Stöhrer, M., et al. Diagnosis and treatment of bladder dysfunction in spinal cord injury patients. *Eur Urol Update Series* 1994. 3: 170.
112. Drake, M., et al., Conservative management in neuropathic urinary incontinence, in *Incontinence*, P. Abrams, L. Cardozo, S. Khoury & A. Wein, Editors. 2013, Health Publication: Plymouth: 2013; pp. 827-1000.
113. Chamberlain, J.D., et al. Mortality and longevity after a spinal cord injury: systematic review and meta-analysis. *Neuroepidemiology*, 2015. 44:182. <http://www.ncbi.nlm.nih.gov/pubmed/25997873>
114. Game, X., et al. Botulinum toxin A detrusor injections in patients with neurogenic detrusor overactivity significantly decrease the incidence of symptomatic urinary tract infections. *Eur Urol*, 2008. 53: 613. <http://www.ncbi.nlm.nih.gov/pubmed/17804150>
115. Frankel, H.L., et al. Long-term survival in spinal cord injury: a fifty year investigation. *Spinal Cord*, 1998. 36: 266. <http://www.ncbi.nlm.nih.gov/pubmed/9589527>
116. Jamil, F. Towards a catheter free status in neurogenic bladder dysfunction: a review of bladder management options in spinal cord injury (SCI). *Spinal Cord*, 2001. 39: 355. <http://www.ncbi.nlm.nih.gov/pubmed/11464308>
117. Thietje, R., et al. Mortality in patients with traumatic spinal cord injury: descriptive analysis of 62 deceased subjects. *J Spinal Cord Med*, 2011. 34: 482. <http://www.ncbi.nlm.nih.gov/pubmed/22118255>



118. Hackler, R.H. A 25-year prospective mortality study in the spinal cord injured patient: comparison with the long-term living paraplegic. *J Urol*, 1977. 117: 486. <http://www.ncbi.nlm.nih.gov/pubmed/850323>
119. Rodrigues, P., et al. Involuntary detrusor contraction is a frequent finding in patients with recurrent urinary tract infections. *Urol Int*, 2014. 93: 67. <http://www.ncbi.nlm.nih.gov/pubmed/25011551>
120. Bauer, S.B. Neurogenic bladder: etiology and assessment. *Pediatr Nephrol*, 2008. 23: 541. <http://www.ncbi.nlm.nih.gov/pubmed/18270749>
121. Barbalias, G.A., et al. Critical evaluation of the Crede maneuver: a urodynamic study of 207 patients. *J Urol*, 1983. 130: 720. <http://www.ncbi.nlm.nih.gov/pubmed/6887405>
122. Reinberg, Y., et al. Renal rupture after the Crede maneuver. *J Pediatr*, 1994. 124: 279. <http://www.ncbi.nlm.nih.gov/pubmed/8301439>
123. Wyndaele, J.J., et al. Neurologic urinary incontinence. *NeuroUrol Urodyn*, 2010. 29: 159. <http://www.ncbi.nlm.nih.gov/pubmed/20025021>
124. Drake, M.J., et al., Neurologic Urinary and Faecal Incontinence., in *Incontinence*, P. Abrams, L. Cardozo, S. Khoury & A. Wein, Editors. 2013, ICUD-EAU.
125. Menon, E.B., et al. Bladder training in patients with spinal cord injury. *Urology*, 1992. 40: 425. <http://www.ncbi.nlm.nih.gov/pubmed/1441039>
126. Furusawa, K., et al. Incidence of symptomatic autonomic dysreflexia varies according to the bowel and bladder management techniques in patients with spinal cord injury. *Spinal Cord*, 2011. 49: 49. <http://www.ncbi.nlm.nih.gov/pubmed/20697419>
127. Outcomes following traumatic spinal cord injury: clinical practice guidelines for health-care professionals. *J Spinal Cord Med*, 2000. 23: 289. <http://www.ncbi.nlm.nih.gov/pubmed/17536300>
128. El-Masri, W.S., et al. Long-term follow-up study of outcomes of bladder management in spinal cord injury patients under the care of the Midlands Centre for Spinal Injuries in Oswestry. *Spinal Cord*, 2012. 50:14. <http://www.ncbi.nlm.nih.gov/pubmed/21808256>
129. Singh, R., et al. Bladder management methods and urological complications in spinal cord injury patients. *Indian J Orthop*, 2011. 45:141. <http://www.ncbi.nlm.nih.gov/pubmed/21430869>
130. Fall, M., et al. Electrical stimulation. A physiologic approach to the treatment of urinary incontinence. *Urol Clin North Am*, 1991. 18: 393. <http://www.ncbi.nlm.nih.gov/pubmed/2017820>
131. Vodusek, D.B., et al. Detrusor inhibition induced by stimulation of pudendal nerve afferents. *NeuroUrol Urodyn*, 1986. 5: 381.
132. Bemelmans, B.L., et al. Neuromodulation by implant for treating lower urinary tract symptoms and dysfunction. *Eur Urol*, 1999. 36: 81. <http://www.ncbi.nlm.nih.gov/pubmed/10420026>
133. Primus, G., et al. Maximal external electrical stimulation for treatment of neurogenic or nonneurogenic urgency and/or urge incontinence. *NeuroUrol Urodyn*, 1996. 15: 187. <http://www.ncbi.nlm.nih.gov/pubmed/8732985>
134. Zecca, C., et al. Maintenance percutaneous posterior nerve stimulation for refractory lower urinary tract symptoms in patients with multiple sclerosis: an open label, multicenter, prospective study. *J Urol*, 2014. 191: 697. <http://www.ncbi.nlm.nih.gov/pubmed/24076308>
135. Kessler, T.M., et al. Transcutaneous electrical nerve stimulation and percutaneous tibial nerve stimulation for neurogenic lower urinary tract dysfunction: a systematic review. *PROSPERO* 2014.
136. McClurg, D., et al. Neuromuscular electrical stimulation and the treatment of lower urinary tract dysfunction in multiple sclerosis-a double blind, placebo controlled, randomised clinical trial. *NeuroUrol Urodyn*, 2008. 27: 231. <http://www.ncbi.nlm.nih.gov/pubmed/17705160>
137. McClurg, D., et al. Comparison of pelvic floor muscle training, electromyography biofeedback, and neuromuscular electrical stimulation for bladder dysfunction in people with multiple sclerosis: a randomized pilot study. *NeuroUrol Urodyn*, 2006. 25: 337. <http://www.ncbi.nlm.nih.gov/pubmed/16637070>
138. Hagerty, J.A., et al. Intravesical electrotherapy for neurogenic bladder dysfunction: a 22-year experience. *J Urol*, 2007. 178: 1680. <http://www.ncbi.nlm.nih.gov/pubmed/17707024>
139. Primus, G., et al. Restoration of micturition in patients with acontractile and hypocontractile detrusor by transurethral electrical bladder stimulation. *NeuroUrol Urodyn*, 1996. 15: 489. <http://www.ncbi.nlm.nih.gov/pubmed/8857617>
140. Lombardi, G., et al. Clinical efficacy of intravesical electrostimulation on incomplete spinal cord patients suffering from chronic neurogenic non-obstructive retention: a 15-year single centre retrospective study. *Spinal Cord*, 2013. 51: 232. <http://www.ncbi.nlm.nih.gov/pubmed/23147136>
141. Brusa, L., et al. Effects of inhibitory rTMS on bladder function in Parkinson's disease patients. *Mov Disord*, 2009. 24: 445. <http://www.ncbi.nlm.nih.gov/pubmed/19133657>
142. Centonze, D., et al. Effects of motor cortex rTMS on lower urinary tract dysfunction in multiple sclerosis. *Mult Scler*, 2007. 13: 269. <http://www.ncbi.nlm.nih.gov/pubmed/17439897>
143. Amend, B., et al. Effective treatment of neurogenic detrusor dysfunction by combined high-dosed antimuscarinics without increased side-effects. *Eur Urol*, 2008. 53: 1021. <http://www.ncbi.nlm.nih.gov/pubmed/18243516>
144. Cameron, A.P. Pharmacologic therapy for the neurogenic bladder. *Urol Clin North Am*, 2010. 37:495. <http://www.ncbi.nlm.nih.gov/pubmed/20955901>
145. Cameron, A.P., et al. Combination drug therapy improves compliance of the neurogenic bladder. *J Urol*, 2009. 182: 1062. <http://www.ncbi.nlm.nih.gov/pubmed/19616807>
146. Thomas, L.H., et al. Treatment of urinary incontinence after stroke in adults. *Cochrane Database Syst Rev*, 2008: CD004462. <http://www.ncbi.nlm.nih.gov/pubmed/18254050>
147. Yeo, L., et al. Urinary tract dysfunction in Parkinson's disease: a review. *Int Urol Nephrol*, 2012. 44:415. <http://www.ncbi.nlm.nih.gov/pubmed/21553114>



148. Stohrer, M., et al. EAU guidelines on neurogenic lower urinary tract dysfunction. *Eur Urol*, 2009. 56: 81. <http://www.ncbi.nlm.nih.gov/pubmed/19403235>
149. Andersson, K.E. Antimuscarinic mechanisms and the overactive detrusor: an update. *Eur Urol*, 2011. 59: 377. <http://www.ncbi.nlm.nih.gov/pubmed/21168951>
150. Kennelly, M.J., et al. Overactive bladder: pharmacologic treatments in the neurogenic population. *Rev Urol*, 2008. 10: 182. <http://www.ncbi.nlm.nih.gov/pubmed/19628264>
151. Madersbacher, H., et al. Neurogenic detrusor overactivity in adults: a review on efficacy, tolerability and safety of oral antimuscarinics. *Spinal Cord*, 2013. 51: 432. <http://www.ncbi.nlm.nih.gov/pubmed/23743498>
152. Madhuvrata, P., et al. Anticholinergic drugs for adult neurogenic detrusor overactivity: a systematic review and meta-analysis. *Eur Urol*, 2012. 62: 816. <http://www.ncbi.nlm.nih.gov/pubmed/22397851>
153. Bennett, N., et al. Can higher doses of oxybutynin improve efficacy in neurogenic bladder? *J Urol*, 2004. 171: 749. <http://www.ncbi.nlm.nih.gov/pubmed/14713802>
154. Horstmann, M., et al. Neurogenic bladder treatment by doubling the recommended antimuscarinic dosage. *Neurourol Urodyn*, 2006. 25: 441. <http://www.ncbi.nlm.nih.gov/pubmed/16847942>
155. Mehnert, U., et al. The management of urinary incontinence in the male neurological patient. *Curr Opin Urol*, 2014. 24: 586. <http://www.ncbi.nlm.nih.gov/pubmed/25389549>
156. Menarini, M., et al. Tropicium chloride in patients with neurogenic detrusor overactivity: is dose titration of benefit to the patients? *Int J Clin Pharmacol Ther*, 2006. 44: 623. <http://www.ncbi.nlm.nih.gov/pubmed/17190372>
157. Nardulli, R., et al. Combined antimuscarinics for treatment of neurogenic overactive bladder. *Int J, Immunopathol Pharmacol*, 2012. 25: 35s. <http://www.ncbi.nlm.nih.gov/pubmed/22652160>
158. Verpoorten, C., et al. The neurogenic bladder: medical treatment. *Pediatr Nephrol*, 2008. 23: 717. <http://www.ncbi.nlm.nih.gov/pubmed/18095004>
159. Isik, A.T., et al. Tropicium and cognition in patients with late onset Alzheimer disease. *J Nutr Health Aging*, 2009. 13: 672. <http://www.ncbi.nlm.nih.gov/pubmed/19657549>
160. Ethans, K.D., et al. Efficacy and safety of tolterodine in people with neurogenic detrusor overactivity. *J Spinal Cord Med*, 2004. 27: 214. <http://www.ncbi.nlm.nih.gov/pubmed/15478523>
161. Mahanta, K., et al. Comparative efficacy and safety of extended-release and instant-release tolterodine in children with neural tube defects having cystometric abnormalities. *J Pediatr Urol*, 2008. 4: 118. <http://www.ncbi.nlm.nih.gov/pubmed/18631906>
162. Reddy, P.P., et al. Long-term efficacy and safety of tolterodine in children with neurogenic detrusor overactivity. *J Pediatr Urol*, 2008. 4: 428. <http://www.ncbi.nlm.nih.gov/pubmed/19013412>
163. Stohrer, M., et al. Propiverine compared to oxybutynin in neurogenic detrusor overactivity-results of a randomized, double-blind, multicenter clinical study. *Eur Urol*, 2007. 51: 235.
164. Madersbacher, H., et al. Efficacy, tolerability and safety profile of propiverine in the treatment of the overactive bladder (non-neurogenic and neurogenic). *World J Urol*, 2001. 19: 324. <http://www.ncbi.nlm.nih.gov/pubmed/11760781>
165. Madersbacher, H., et al. Propiverine vs oxybutynin for treating neurogenic detrusor overactivity in children and adolescents: results of a multicentre observational cohort study. *BJU Int*, 2009. 103: 776. <http://www.ncbi.nlm.nih.gov/pubmed/19007380>
166. Stohrer, M., et al. Efficacy and tolerability of propiverine hydrochloride extended-release compared with immediate-release in patients with neurogenic detrusor overactivity. *Spinal Cord*, 2013. 51: 419. <http://www.ncbi.nlm.nih.gov/pubmed/23338657>
167. Nicholas, R.S., et al. Anticholinergics for urinary symptoms in multiple sclerosis. *Cochrane Database Syst Rev*, 2009: CD004193. <http://www.ncbi.nlm.nih.gov/pubmed/19160231>
168. van Rey, F., et al. Solifenacin in multiple sclerosis patients with overactive bladder: a prospective study. *Adv Urol*, 2011. 2011: 834753. <http://www.ncbi.nlm.nih.gov/pubmed/21687581>
169. Carl, S., et al. Darifenacin is also effective in neurogenic bladder dysfunction (multiple sclerosis). *Urology*, 2006. 68: 250.
170. Bycroft, J., et al. The effect of darifenacin on neurogenic detrusor overactivity in patients with spinal cord injury. *Neurourol Urodyn* 2003. 22: A190. <http://www.ics.org/Abstracts/Publish/41/000190.pdf>
171. Hassouna, M. Comparative Study of the Efficacy and Safety of Muscarinic M3 Receptors Antagonists in the Treatment of Neurogenic Detrusor Overactivity (NCT00800462). <http://clinicaltrials.gov/show/NCT00800462>
172. Astellas Pharma Inc. Clinical study of solifenacin succinate in patients with bladder symptoms due to spinal cord injury or multiple sclerosis (SONIC). (NCT00629642). January 2011 (final data collection date for primary outcome measure). <http://clinicaltrials.gov/show/NCT00629642>
173. Zesiewicz, T., et al., Solifenacin succinate (VESicare) for the treatment of overactive bladder in Parkinson's Disease (URGE-PD). in University of South Florida August 2014 (final data collection date for primary outcome measure): NCT01018264. <https://www.clinicaltrials.gov/show/NCT01018264>
174. Appell, R.A. Pharmacotherapy for overactive bladder: an evidence-based approach to selecting an antimuscarinic agent. *Drugs*, 2006. 66: 1361. <http://www.ncbi.nlm.nih.gov/pubmed/16903770>
175. Kennelly, M.J., et al. Efficacy and safety of oxybutynin transdermal system in spinal cord injury patients with neurogenic detrusor overactivity and incontinence: an open-label, dose-titration study. *Urology*, 2009. 74: 741. <http://www.ncbi.nlm.nih.gov/pubmed/19628264>
176. Wöllner, J., et al. Initial experience with the treatment of neurogenic detrusor overactivity with a new beta-3 agonist (mirabegron) in patients with spinal cord injury. *Spinal Cord*, 2015. <http://www.ncbi.nlm.nih.gov/pubmed/26503222>
177. Welk, B., Urodynamic and Clinical Efficacy of Mirabegron for Neurogenic Bladder Patients, Ongoing study: ClinicalTrials.gov Identifier: NCT02044510.



178. Mariano, P.S., et al. Evidence for a pH-dependent irreversible formation of a stable conformation of phenacyl-alpha-chymotrypsin. *Biochem J*, 1978. 171: 115. <http://www.ncbi.nlm.nih.gov/pubmed/25656>
179. Barendrecht, M.M., et al. Is the use of parasympathomimetics for treating an underactive urinary bladder evidence-based? *BJU Int*, 2007. 99: 749. <http://www.ncbi.nlm.nih.gov/pubmed/17233798>
180. Apostolidis, A. Taming the cannabinoids: new potential in the pharmacologic control of lower urinary tract dysfunction. *Eur Urol*, 2012. 61:107. <http://www.ncbi.nlm.nih.gov/pubmed/21996529>
181. Gratzke, C., et al. Effects of cannabimimetic, a novel selective cannabinoid 2 receptor agonist, on bladder function in normal rats. *Eur Urol*, 2010. 57:1093. <http://www.ncbi.nlm.nih.gov/pubmed/20207474>
182. Kavia, R.B., et al. Randomized controlled trial of Sativex to treat detrusor overactivity in multiple sclerosis. *Mult Scler*, 2010. 16: 1349. <http://www.ncbi.nlm.nih.gov/pubmed/20829244>
183. Abrams, P., et al. Tamsulosin: efficacy and safety in patients with neurogenic lower urinary tract dysfunction due to suprasacral spinal cord injury. *J Urol*, 2003. 170: 1242. <http://www.ncbi.nlm.nih.gov/pubmed/14501734>
184. Guttmann, L., et al. The value of intermittent catheterisation in the early management of traumatic paraplegia and tetraplegia. *Paraplegia*, 1966. 4: 63. <http://www.ncbi.nlm.nih.gov/pubmed/5969402>
185. Lapides, J., et al. Clean, intermittent self-catheterization in the treatment of urinary tract disease. *J Urol*, 1972. 107: 458. <http://www.ncbi.nlm.nih.gov/pubmed/5010715>
186. Wyndaele, J.J. Intermittent catheterization: which is the optimal technique? *Spinal Cord*, 2002. 40:432. <http://www.ncbi.nlm.nih.gov/pubmed/12185603>
187. Prieto-Fingerhut, T., et al. A study comparing sterile and nonsterile urethral catheterization in patients with spinal cord injury. *Rehabil Nurs*, 1997. 22: 299. <http://www.ncbi.nlm.nih.gov/pubmed/9416190>
188. Kiddoo, D., et al. Randomized Crossover Trial of Single Use Hydrophilic Coated vs Multiple Use Polyvinylchloride Catheters for Intermittent Catheterization to Determine Incidence of Urinary Infection. *J Urol*, 2015. 194: 174. <http://www.ncbi.nlm.nih.gov/pubmed/25584995>
189. Günther, M., et al. Auswirkungen des aseptischen intermittierenden Katheterismus auf die männliche Harnröhre. *Der Urologe B*, 2001. 41: 359. <http://link.springer.com/article/10.1007%2Fs001310170044>
190. Bakke, A., et al. Physical predictors of infection in patients treated with clean intermittent catheterization: a prospective 7-year study. *Br J Urol*, 1997. 79: 85. <http://www.ncbi.nlm.nih.gov/pubmed/9043503>
191. Waller, L., et al. Clean intermittent catheterization in spinal cord injury patients: long-term followup of a hydrophilic low friction technique. *J Urol*, 1995. 153: 345. <http://www.ncbi.nlm.nih.gov/pubmed/7815579>
192. Wyndaele, J.J. Complications of intermittent catheterization: their prevention and treatment. *Spinal Cord*, 2002. 40: 536. <http://www.ncbi.nlm.nih.gov/pubmed/12235537>
193. Woodbury, M.G., et al. Intermittent catheterization practices following spinal cord injury: a national survey. *Can J Urol*, 2008. 15: 4065. <http://www.ncbi.nlm.nih.gov/pubmed/18570710>
194. Weld, K.J., et al. Effect of bladder management on urological complications in spinal cord injured patients. *J Urol*, 2000. 163: 768. <http://www.ncbi.nlm.nih.gov/pubmed/10687973>
195. Bennett, C.J., et al. Comparison of bladder management complication outcomes in female spinal cord injury patients. *J Urol*, 1995. 153: 1458. <http://www.ncbi.nlm.nih.gov/pubmed/7714965>
196. Chancellor, M.B., et al. Functional urethral closure with pubovaginal sling for destroyed female urethra after long-term urethral catheterization. *Urology*, 1994. 43: 499. <http://www.ncbi.nlm.nih.gov/pubmed/8154071>
197. Chao, R., et al. Fate of upper urinary tracts in patients with indwelling catheters after spinal cord injury. *Urology*, 1993. 42: 259. <http://www.ncbi.nlm.nih.gov/pubmed/8379025>
198. Larsen, L.D., et al. Retrospective analysis of urologic complications in male patients with spinal cord injury managed with and without indwelling urinary catheters. *Urology*, 1997. 50: 418. <http://www.ncbi.nlm.nih.gov/pubmed/9301708>
199. Mitsui, T., et al. Is suprapubic cystostomy an optimal urinary management in high quadriplegics?. A comparative study of suprapubic cystostomy and clean intermittent catheterization. *Eur Urol*, 2000. 38: 434. <http://www.ncbi.nlm.nih.gov/pubmed/11025382>
200. Park, Y.I., et al. A method to minimize indwelling catheter calcification and bladder stones in individuals with spinal cord injury. *J Spinal Cord Med*, 2001. 24: 105. <http://www.ncbi.nlm.nih.gov/pubmed/11587416>
201. Weld, K.J., et al. Influences on renal function in chronic spinal cord injured patients. *J Urol*, 2000. 164: 1490. <http://www.ncbi.nlm.nih.gov/pubmed/11025689>
202. West, D.A., et al. Role of chronic catheterization in the development of bladder cancer in patients with spinal cord injury. *Urology*, 1999. 53: 292. <http://www.ncbi.nlm.nih.gov/pubmed/9933042>
203. Hollingsworth, J.M., et al. Determining the noninfectious complications of indwelling urethral catheters: a systematic review and meta-analysis. *Ann Intern Med*, 2013. 159: 401. <http://www.ncbi.nlm.nih.gov/pubmed/24042368>
204. Di Stasi, S.M., et al. Intravesical oxybutynin: mode of action assessed by passive diffusion and electromotive administration with pharmacokinetics of oxybutynin and N-desethyl oxybutynin. *J Urol*, 2001. 166: 2232. <http://www.ncbi.nlm.nih.gov/pubmed/11696741>
205. Buyse, G., et al. Intravesical oxybutynin for neurogenic bladder dysfunction: less systemic side effects due to reduced first pass metabolism. *J Urol*, 1998. 160: 892. <http://www.ncbi.nlm.nih.gov/pubmed/9720583>
206. Haferkamp, A., et al. Dosage escalation of intravesical oxybutynin in the treatment of neurogenic bladder patients. *Spinal Cord*, 2000. 38: 250. <http://www.ncbi.nlm.nih.gov/pubmed/10822396>



207. Pannek, J., et al. Combined intravesical and oral oxybutynin chloride in adult patients with spinal cord injury. *Urology*, 2000. 55: 358. <http://www.ncbi.nlm.nih.gov/pubmed/10699610>
208. Schroder, A., et al. Efficacy, safety, and tolerability of intravesically administered 0.1% oxybutynin hydrochloride solution in adult patients with neurogenic bladder: A randomized, prospective, controlled multi-center trial. *NeuroUrol Urodyn*, 2015. <http://www.ncbi.nlm.nih.gov/pubmed/25754454>
209. Geirsson, G., et al. Clinical and urodynamic effects of intravesical capsaicin treatment in patients with chronic traumatic spinal detrusor hyperreflexia. *J Urol*, 1995. 154: 1825. <http://www.ncbi.nlm.nih.gov/pubmed/7563356>
210. Giannantoni, A., et al. Intravesical resiniferatoxin versus botulinum-A toxin injections for neurogenic detrusor overactivity: a prospective randomized study. *J Urol*, 2004. 172: 240. <http://www.ncbi.nlm.nih.gov/pubmed/15201783>
211. Kim, J.H., et al. Intravesical resiniferatoxin for refractory detrusor hyperreflexia: a multicenter, blinded, randomized, placebo-controlled trial. *J Spinal Cord Med*, 2003. 26: 358. <http://www.ncbi.nlm.nih.gov/pubmed/14992337>
212. Del Popolo, G., et al. Neurogenic detrusor overactivity treated with english botulinum toxin a: 8-year experience of one single centre. *Eur Urol*, 2008. 53: 1013. <http://www.ncbi.nlm.nih.gov/pubmed/17950989>
213. Reitz, A., et al. European experience of 200 cases treated with botulinum-A toxin injections into the detrusor muscle for urinary incontinence due to neurogenic detrusor overactivity. *Eur Urol*, 2004. 45: 510. <http://www.ncbi.nlm.nih.gov/pubmed/15041117>
214. Schurch, B., et al. Botulinum toxin type a is a safe and effective treatment for neurogenic urinary incontinence: results of a single treatment, randomized, placebo controlled 6-month study. *J Urol*, 2005. 174: 196. <http://www.ncbi.nlm.nih.gov/pubmed/15947626>
215. Cruz, F., et al. Efficacy and safety of onabotulinumtoxinA in patients with urinary incontinence due to neurogenic detrusor overactivity: a randomised, double-blind, placebo-controlled trial. *Eur Urol*, 2011. 60: 742. <http://www.ncbi.nlm.nih.gov/pubmed/21798658>
216. Ginsberg, D., et al. Phase 3 efficacy and tolerability study of onabotulinumtoxinA for urinary incontinence from neurogenic detrusor overactivity. *J Urol*, 2012. 187: 2131. <http://www.ncbi.nlm.nih.gov/pubmed/22503020>
217. Mehta, S., et al. Meta-analysis of botulinum toxin A detrusor injections in the treatment of neurogenic detrusor overactivity after spinal cord injury. *Arch Phys Med Rehabil*, 2013. 94: 1473. <http://www.ncbi.nlm.nih.gov/pubmed/23632286>
218. Mangera, A., et al. An updated systematic review and statistical comparison of standardised mean outcomes for the use of botulinum toxin in the management of lower urinary tract disorders. *Eur Urol*, 2014. 65: 981. <http://www.ncbi.nlm.nih.gov/pubmed/24239446>
219. Grosse, J., et al. Success of repeat detrusor injections of botulinum a toxin in patients with severe neurogenic detrusor overactivity and incontinence. *Eur Urol*, 2005. 47: 653. <http://www.ncbi.nlm.nih.gov/pubmed/15826758>
220. Dykstra, D.D., et al. Treatment of detrusor-sphincter dyssynergia with botulinum A toxin: a doubleblind study. *Arch Phys Med Rehabil*, 1990. 71: 24. <http://www.ncbi.nlm.nih.gov/pubmed/2297305>
221. Petit, H., et al. Botulinum A toxin treatment for detrusor-sphincter dyssynergia in spinal cord disease. *Spinal Cord*, 1998. 36: 91. <http://www.ncbi.nlm.nih.gov/pubmed/9494997>
222. Schurch, B., et al. Botulinum-A toxin as a treatment of detrusor-sphincter dyssynergia: a prospective study in 24 spinal cord injury patients. *J Urol*, 1996. 155: 1023. <http://www.ncbi.nlm.nih.gov/pubmed/8583552>
223. Utomo, E., et al. Surgical management of functional bladder outlet obstruction in adults with neurogenic bladder dysfunction. *Cochrane Database Syst Rev*, 2014. 5: CD004927. <http://www.ncbi.nlm.nih.gov/pubmed/24859260>
224. Chancellor, M.B., et al. Prospective comparison of external sphincter balloon dilatation and prosthesis placement with external sphincterotomy in spinal cord injured men. *Arch Phys Med Rehabil*, 1994. 75: 297. <http://www.ncbi.nlm.nih.gov/pubmed/8129583>
225. Perkasch, I. Use of contact laser crystal tip firing Nd:YAG to relieve urinary outflow obstruction in male neurogenic bladder patients. *J Clin Laser Med Surg*, 1998. 16: 33. <http://www.ncbi.nlm.nih.gov/pubmed/9728128>
226. Reynard, J.M., et al. Sphincterotomy and the treatment of detrusor-sphincter dyssynergia: current status, future prospects. *Spinal Cord*, 2003. 41: 1. <http://www.ncbi.nlm.nih.gov/pubmed/12494314>
227. Noll, F., et al. Transurethral sphincterotomy in quadriplegic patients: long-term-follow-up. *NeuroUrol Urodyn*, 1995. 14: 351. <http://www.ncbi.nlm.nih.gov/pubmed/7581471>
228. Derry, F., et al. Audit of bladder neck resection in spinal cord injured patients. *Spinal Cord*, 1998. 36: 345. <http://www.ncbi.nlm.nih.gov/pubmed/9601115>
229. Chancellor, M.B., et al. Long-term followup of the North American multicenter UroLume trial for the treatment of external detrusor-sphincter dyssynergia. *J Urol*, 1999. 161: 1545. <http://www.ncbi.nlm.nih.gov/pubmed/10210393>
230. Seoane-Rodriguez, S., et al. Long-term follow-up study of intraurethral stents in spinal cord injured patients with detrusor-sphincter dyssynergia. *Spinal Cord*, 2007. 45: 621. <http://www.ncbi.nlm.nih.gov/pubmed/17211463>
231. Gajewski, J.B., et al. Removal of UroLume endoprosthesis: experience of the North American Study Group for detrusor-sphincter dyssynergia application. *J Urol*, 2000. 163: 773. <http://www.ncbi.nlm.nih.gov/pubmed/10687974>
232. Wilson, T.S., et al. UroLume stents: lessons learned. *J Urol*, 2002. 167: 2477. <http://www.ncbi.nlm.nih.gov/pubmed/11992061>
233. Polguer, T., et al. [Treatment of detrusor-striated sphincter dyssynergia with permanent nitinol urethral stent: results after a minimum follow-up of 2 years]. *Prog Urol*, 2012. 22: 1058. <http://www.ncbi.nlm.nih.gov/pubmed/23182120>
234. van der Merwe, A., et al. Outcome of dual flange metallic urethral stents in the treatment of neuropathic bladder dysfunction after spinal cord injury. *J Endourol*, 2012. 26: 1210. <http://www.ncbi.nlm.nih.gov/pubmed/22519741>





235. Pannek, J., et al. Clinical usefulness of the memokath stent as a second-line procedure after sphincterotomy failure. *J Endourol*, 2011. 25: 335. <http://www.ncbi.nlm.nih.gov/pubmed/20977372>
236. Abdul-Rahman, A., et al. A 20-year follow-up of the mesh wallstent in the treatment of detrusor external sphincter dyssynergia in patients with spinal cord injury. *BJU Int*, 2010. 106: 1510. <http://www.ncbi.nlm.nih.gov/pubmed/20500511>
237. Bennett, J.K., et al. Collagen injections for intrinsic sphincter deficiency in the neuropathic urethra. *Paraplegia*, 1995. 33: 697. <http://www.ncbi.nlm.nih.gov/pubmed/8927407>
238. Block, C.A., et al. Long-term efficacy of periurethral collagen injection for the treatment of urinary incontinence secondary to myelomeningocele. *J Urol*, 2003. 169: 327. <http://www.ncbi.nlm.nih.gov/pubmed/12478183>
239. Schurch, B., et al. Intraurethral sphincter prosthesis to treat hyporeflexic bladders in women: does it work? *BJU Int*, 1999. 84: 789. <http://www.ncbi.nlm.nih.gov/pubmed/10532973>
240. Barthold, J.S., et al. Results of the rectus fascial sling and wrap procedures for the treatment of neurogenic sphincteric incontinence. *J Urol*, 1999. 161: 272. <http://www.ncbi.nlm.nih.gov/pubmed/10037423>
241. Daneshmand, S., et al. Puboprosthetic sling repair for treatment of urethral incompetence in adult neurogenic incontinence. *J Urol*, 2003. 169: 199. <http://www.ncbi.nlm.nih.gov/pubmed/12478135>
242. Gormley, E.A., et al. Pubovaginal slings for the management of urinary incontinence in female adolescents. *J Urol*, 1994. 152: 822. <http://www.ncbi.nlm.nih.gov/pubmed/8022024>
243. Herschorn, S., et al. Fascial slings and bladder neck tapering in the treatment of male neurogenic incontinence. *J Urol*, 1992. 147: 1073. <http://www.ncbi.nlm.nih.gov/pubmed/1552586>
244. Kakizaki, H., et al. Fascial sling for the management of urinary incontinence due to sphincter incompetence. *J Urol*, 1995. 153: 644. <http://www.ncbi.nlm.nih.gov/pubmed/7861504>
245. Mingin, G.C., et al. The rectus myofascial wrap in the management of urethral sphincter incompetence. *BJU Int*, 2002. 90: 550. <http://www.ncbi.nlm.nih.gov/pubmed/12230615>
246. Abdul-Rahman, A., et al. Long-term outcome of tension-free vaginal tape for treating stress incontinence in women with neuropathic bladders. *BJU Int*, 2010. 106: 827. <http://www.ncbi.nlm.nih.gov/pubmed/20132201>
247. Losco, G.S., et al. Long-term outcome of transobturator tape (TOT) for treatment of stress urinary incontinence in females with neuropathic bladders. *Spinal Cord*, 2015. 53: 544. <http://www.ncbi.nlm.nih.gov/pubmed/25917951>
248. Athanasopoulos, A., et al. Treating stress urinary incontinence in female patients with neuropathic bladder: the value of the autologous fascia rectus sling. *Int Urol Nephrol*, 2012. 44: 1363. <http://www.ncbi.nlm.nih.gov/pubmed/22821050>
249. Groen, L.A., et al. The AdVance male sling as a minimally invasive treatment for intrinsic sphincter deficiency in patients with neurogenic bladder sphincter dysfunction: a pilot study. *Neurourol Urodyn*, 2012. 31: 1284. <http://www.ncbi.nlm.nih.gov/pubmed/22847896>
250. Mehnert, U., et al. Treatment of neurogenic stress urinary incontinence using an adjustable continence device: 4-year followup. *J Urol*, 2012. 188: 2274. <http://www.ncbi.nlm.nih.gov/pubmed/23083648>
251. Light, J.K., et al. Use of the artificial urinary sphincter in spinal cord injury patients. *J Urol*, 1983. 130: 1127. <http://www.ncbi.nlm.nih.gov/pubmed/6644893>
252. Costa, P., et al. Long-term results of artificial urinary sphincter for women with type III stress urinary incontinence. *Eur Urol*, 2013. 63: 753. <http://www.ncbi.nlm.nih.gov/pubmed/22445222>
253. Thomas, K., et al. Outcome of the artificial urinary sphincter in female patients. *J Urol*, 2002. 167: 1720. <http://www.ncbi.nlm.nih.gov/pubmed/11912395>
254. Chartier Kastler, E., et al. Treatment of neurogenic male urinary incontinence related to intrinsic sphincter insufficiency with an artificial urinary sphincter: a French retrospective multicentre study. *BJU Int*, 2011. 107: 426. <http://www.ncbi.nlm.nih.gov/pubmed/20633005>
255. Elliott, D.S., et al. Mayo Clinic long-term analysis of the functional durability of the AMS 800 artificial urinary sphincter: a review of 323 cases. *J Urol*, 1998. 159: 1206. <http://www.ncbi.nlm.nih.gov/pubmed/9507835>
256. Fulford, S.C., et al. The fate of the 'modern' artificial urinary sphincter with a follow-up of more than 10 years. *Br J Urol*, 1997. 79: 713. <http://www.ncbi.nlm.nih.gov/pubmed/9158507>
257. Viers, B.R., et al. Simultaneous augmentation cystoplasty and cuff only artificial urinary sphincter in children and young adults with neurogenic urinary incontinence. *J Urol*, 2014. 191: 1104. <http://www.ncbi.nlm.nih.gov/pubmed/24060640>
258. Janknegt, R.A., et al. Electrically stimulated gracilis sphincter for treatment of bladder sphincter incontinence. *Lancet*, 1992. 340: 1129. <http://www.ncbi.nlm.nih.gov/pubmed/1359213>
259. Chancellor, M.B., et al. Gracilis muscle transposition with electrical stimulation for sphincteric incontinence: a new approach. *World J Urol*, 1997. 15: 320. <http://www.ncbi.nlm.nih.gov/pubmed/9372585>
260. Chancellor, M.B., et al. Gracilis urethromyoplasty--an autologous urinary sphincter for neurologically impaired patients with stress incontinence. *Spinal Cord*, 1997. 35: 546. <http://www.nature.com/sc/journal/v35/n8/abs/3100444a.html>
261. Donnahoo, K.K., et al. The Young-Dees-Leadbetter bladder neck repair for neurogenic incontinence. *J Urol*, 1999. 161: 1946. <http://www.ncbi.nlm.nih.gov/pubmed/10332478>
262. Kropp, K.A., et al. Urethral lengthening and reimplantation for neurogenic incontinence in children. *J Urol*, 1986. 135: 533. <http://www.ncbi.nlm.nih.gov/pubmed/3944902>
263. Salle, J.L., et al. Urethral lengthening with anterior bladder wall flap (Pippi Salle procedure): modifications and extended indications of the technique. *J Urol*, 1997. 158: 585. <http://www.ncbi.nlm.nih.gov/pubmed/9224369>
264. Rawashdeh, Y.F., et al. International Children's Continence Society's recommendations for therapeutic intervention in congenital neuropathic bladder and bowel dysfunction in children. *Neurourol Urodyn*, 2012. 31: 615. <http://www.ncbi.nlm.nih.gov/pubmed/22532368>



265. Nagib, A., et al. Successful control of selective anterior sacral rhizotomy for treatment of spastic bladder and ureteric reflux in paraplegics. *Med Serv J Can*, 1966. 22: 576. <http://www.ncbi.nlm.nih.gov/pubmed/5966992>
266. Schneidau, T., et al. Selective sacral rhizotomy for the management of neurogenic bladders in spina bifida patients: long-term followup. *J Urol*, 1995. 154: 766. <http://www.ncbi.nlm.nih.gov/pubmed/7609174>
267. Young, B., et al. Percutaneous sacral rhizotomy for neurogenic detrusor hyperreflexia. *J Neurosurg*, 1980. 53: 85. <http://www.ncbi.nlm.nih.gov/pubmed/7411212>
268. Koldewijn, E.L., et al. Bladder compliance after posterior sacral root rhizotomies and anterior sacral root stimulation. *J Urol*, 1994. 151: 955. <http://www.ncbi.nlm.nih.gov/pubmed/8126835>
269. Krasnik, D., et al. Urodynamic results, clinical efficacy, and complication rates of sacral intradural deafferentation and sacral anterior root stimulation in patients with neurogenic lower urinary tract dysfunction resulting from complete spinal cord injury. *Neurourol Urodyn*, 2014. 33: 1202. <http://www.ncbi.nlm.nih.gov/pubmed/24038405>
270. Singh, G., et al. Intravesical oxybutynin in patients with posterior rhizotomies and sacral anterior root stimulators. *Neurourol Urodyn*, 1995. 14: 65. <http://www.ncbi.nlm.nih.gov/pubmed/7742851>
271. Van Kerrebroeck, P.E., et al. Results of the treatment of neurogenic bladder dysfunction in spinal cord injury by sacral posterior root rhizotomy and anterior sacral root stimulation. *J Urol*, 1996. 155:1378. <http://www.ncbi.nlm.nih.gov/pubmed/8632580>
272. Kutzenberger, J.S. Surgical therapy of neurogenic detrusor overactivity (hyperreflexia) in paraplegic patients by sacral deafferentation and implant driven micturition by sacral anterior root stimulation: methods, indications, results, complications, and future prospects. *Acta Neurochir*, 2007. 97: 333. <http://www.ncbi.nlm.nih.gov/pubmed/17691394>
273. Bhadra, N., et al. Selective suppression of sphincter activation during sacral anterior nerve root stimulation. *Neurourol Urodyn*, 2002. 21: 55. <http://www.ncbi.nlm.nih.gov/pubmed/11835425>
274. Kirkham, A.P., et al. Neuromodulation through sacral nerve roots 2 to 4 with a Finetech-Brindley sacral posterior and anterior root stimulator. *Spinal Cord*, 2002. 40: 272. <http://www.ncbi.nlm.nih.gov/pubmed/12037708>
275. Schumacher, S., et al. Extradural cold block for selective neurostimulation of the bladder: development of a new technique. *J Urol*, 1999. 161: 950. <http://www.ncbi.nlm.nih.gov/pubmed/10022732>
276. Brindley, G.S. An implant to empty the bladder or close the urethra. *J Neurol Neurosurg Psychiatry*, 1977. 40: 358. <http://www.ncbi.nlm.nih.gov/pubmed/406364>
277. Benard, A., et al. Comparative cost-effectiveness analysis of sacral anterior root stimulation for rehabilitation of bladder dysfunction in spinal cord injured patients. *Neurosurgery*, 2013. 73: 600. <http://www.ncbi.nlm.nih.gov/pubmed/23787880>
278. Martens, F.M., et al. Quality of life in complete spinal cord injury patients with a Brindley bladder stimulator compared to a matched control group. *Neurourol Urodyn*, 2011. 30: 551. <http://www.ncbi.nlm.nih.gov/pubmed/21328472>
279. Wollner, J., et al. Surgery Illustrated - surgical atlas sacral neuromodulation. *BJU Int*, 2012. 110: 146. <http://www.ncbi.nlm.nih.gov/pubmed/22691023>
280. Kessler, T.M., et al. Sacral neuromodulation for neurogenic lower urinary tract dysfunction: systematic review and meta-analysis. *Eur Urol*, 2010. 58: 865. <http://www.ncbi.nlm.nih.gov/pubmed/20934242>
281. Lombardi, G., et al. Sacral neuromodulation for neurogenic non-obstructive urinary retention in incomplete spinal cord patients: a ten year follow-up single-centre experience. *Spinal Cord*, 2014. 52: 241. <http://www.ncbi.nlm.nih.gov/pubmed/24394604>
282. Lay, A.H., et al. The role of neuromodulation in patients with neurogenic overactive bladder. *Curr Urol Rep*, 2012. 13: 343. <http://www.ncbi.nlm.nih.gov/pubmed/22865208>
283. Zhang, Y.H., et al. Enveloping the bladder with displacement of flap of the rectus abdominis muscle for the treatment of neurogenic bladder. *J Urol*, 1990. 144: 1194. <http://www.ncbi.nlm.nih.gov/pubmed/2146404>
284. Stenzl, A., et al. Restoration of voluntary emptying of the bladder by transplantation of innervated free skeletal muscle. *Lancet*, 1998. 351:1483. <http://www.ncbi.nlm.nih.gov/pubmed/9605805>
285. Gakis, G., et al. Functional detrusor myoplasty for bladder acontractility: long-term results. *J Urol*, 2011. 185: 593. <http://www.ncbi.nlm.nih.gov/pubmed/2116886>
286. Ninkovic, M., et al. The latissimus dorsi detrusor myoplasty for functional treatment of bladder acontractility. *Clin Plast Surg*, 2012. 39:507. <http://www.ncbi.nlm.nih.gov/pubmed/23036300>
287. Braren, V., et al. Laparoscopic bladder autoaugmentation in children. *Urol Clin North Am*, 1998. 25:533. <http://www.ncbi.nlm.nih.gov/pubmed/9728222>
288. Cartwright, P.C., et al. Bladder autoaugmentation: early clinical experience. *J Urol*, 1989. 142: 505. <http://www.ncbi.nlm.nih.gov/pubmed/2746767>
289. Duel, B.P., et al. Alternative techniques for augmentation cystoplasty. *J Urol*, 1998. 159: 998. <http://www.ncbi.nlm.nih.gov/pubmed/9474216>
290. Poppas, D.P., et al. Laparoscopic laser assisted autoaugmentation of the pediatric neurogenic bladder: early experience with urodynamic followup. *J Urol*, 1996. 155: 1057. <http://www.ncbi.nlm.nih.gov/pubmed/8583564>
291. Snow, B.W., et al. Bladder autoaugmentation. *Urol Clin North Am*, 1996. 23: 323. <http://www.ncbi.nlm.nih.gov/pubmed/8659030>
292. Stohrer, M., et al. Bladder auto-augmentation--an alternative for enterocystoplasty: preliminary results. *Neurourol Urodyn*, 1995. 14:11. <http://www.ncbi.nlm.nih.gov/pubmed/7742844>
293. Stohrer, M., et al. Bladder autoaugmentation in adult patients with neurogenic voiding dysfunction. *Spinal Cord*, 1997. 35: 456. <http://www.ncbi.nlm.nih.gov/pubmed/9232751>
294. Vainrib, M., et al. Differences in urodynamic study variables in adult patients with neurogenic bladder and myelomeningocele before and after augmentation enterocystoplasty. *Neurourol Urodyn*, 2013. 32: 250. <http://www.ncbi.nlm.nih.gov/pubmed/22965686>



295. Krebs, J., et al. Functional outcome of supratrigonal cystectomy and augmentation ileocystoplasty in adult patients with refractory neurogenic lower urinary tract dysfunction. *Neurourol Urodyn*, 2014. <http://www.ncbi.nlm.nih.gov/pubmed/25524480>
296. Gough, D.C. Enterocystoplasty. *BJU Int*, 2001. 88: 739. <http://www.ncbi.nlm.nih.gov/pubmed/11890246>
297. Greenwell, T.J., et al. Augmentation cystoplasty. *BJU Int*, 2001. 88: 511. <http://www.ncbi.nlm.nih.gov/pubmed/11678743>
298. Vajda, P., et al. Histological findings after colocolocystoplasty and gastrocystoplasty. *J Urol*, 2002. 168:698. <http://www.ncbi.nlm.nih.gov/pubmed/12131353>
299. Chapple, C.R., et al. Surgery for detrusor overactivity. *World J Urol*, 1998. 16: 268. <http://www.ncbi.nlm.nih.gov/pubmed/9775426>
300. Comer, M.T., et al. Reconstruction of the urinary bladder by auto-augmentation, enterocystoplasty, and composite enterocystoplasty. *Adv Exp Med Biol*, 1999. 462: 43. <http://www.ncbi.nlm.nih.gov/pubmed/10599412>
301. Cranidis, A., et al. Bladder augmentation. *Int Urogynecol J Pelvic Floor Dysfunct*, 2000. 11: 33. <http://www.ncbi.nlm.nih.gov/pubmed/10738932>
302. Leng, W.W., et al. Enterocystoplasty or detrusor myectomy? Comparison of indications and outcomes for bladder augmentation. *J Urol*, 1999. 161:758. <http://www.ncbi.nlm.nih.gov/pubmed/10022679>
303. Niknejad, K.G., et al. Bladder augmentation techniques in women. *Int Urogynecol J Pelvic Floor Dysfunct*, 2000. 11: 156. <http://www.ncbi.nlm.nih.gov/pubmed/11484743>
304. Oge, O., et al. Urothelium-preserving augmentation cystoplasty covered with a peritoneal flap. *BJU Int*, 2000. 85: 802. <http://www.ncbi.nlm.nih.gov/pubmed/10792156>
305. Siracusano, S., et al. Laparoscopic bladder auto-augmentation in an incomplete traumatic spinal cord injury. *Spinal Cord*, 2000. 38: 59. <http://www.ncbi.nlm.nih.gov/pubmed/10762200>
306. Westney, O.L., et al. Surgical procedures for the treatment of urge incontinence. *Tech Urol*, 2001. 7:126. <http://www.ncbi.nlm.nih.gov/pubmed/11383990>
307. Zhang, F., et al. Sigmoidocolocystoplasty with ureteral reimplantation for treatment of neurogenic bladder. *Urology*, 2012. 80: 440. <http://www.ncbi.nlm.nih.gov/pubmed/22857763>
308. Khavari, R., et al. A modification to augmentation cystoplasty with catheterizable stoma for neurogenic patients: technique and long-term results. *Urology*, 2012. 80: 460. <http://www.ncbi.nlm.nih.gov/pubmed/22704181>
309. Hadley, D., et al. Creation of a continent urinary channel in adults with neurogenic bladder: long term results with the Monti and Casale (Spiral Monti) procedures. *Urology*, 2014. 83: 1176. <http://www.ncbi.nlm.nih.gov/pubmed/24612618>
310. Leslie, B., et al. Long-term followup and time to event outcome analysis of continent catheterizable channels. *J Urol*, 2011. 185: 2298. <http://www.ncbi.nlm.nih.gov/pubmed/21511280>
311. Duckett, J.W., et al. Appendicovesicostomy (and variations) in bladder reconstruction. *J Urol*, 1993. 149: 567. <http://www.ncbi.nlm.nih.gov/pubmed/8437267>
312. Kajbafzadeh, A.M., et al. Simultaneous Malone antegrade continent enema and Mitrofanoff principle using the divided appendix: report of a new technique for prevention of stoma complications. *J Urol*, 2001. 165: 2404. <http://www.ncbi.nlm.nih.gov/pubmed/11371987>
313. Kawai, K., et al. Tissue-engineered artificial urothelium. *World J Surg*, 2000. 24: 1160. <http://www.ncbi.nlm.nih.gov/pubmed/11071451>
314. Liard, A., et al. The Mitrofanoff procedure: 20 years later. *J Urol*, 2001. 165: 2394. <http://www.ncbi.nlm.nih.gov/pubmed/11371985>
315. Moreno, J.G., et al. Improved quality of life and sexuality with continent urinary diversion in quadriplegic women with umbilical stoma. *Arch Phys Med Rehabil*, 1995. 76: 758. <http://www.ncbi.nlm.nih.gov/pubmed/7632132>
316. Sekar, P., et al. Comparison of long-term renal function after spinal cord injury using different urinary management methods. *Arch Phys Med Rehabil*, 1997. 78: 992. <http://www.ncbi.nlm.nih.gov/pubmed/9305274>
317. Stein, R., et al. Urinary diversion and orthotopic bladder substitution in children and young adults with neurogenic bladder: a safe option for treatment? *J Urol*, 2000. 163: 568. <http://www.ncbi.nlm.nih.gov/pubmed/1064768>
318. Sylora, J.A., et al. Intermittent self-catheterization by quadriplegic patients via a catheterizable Mitrofanoff channel. *J Urol*, 1997. 157: 48. <http://www.ncbi.nlm.nih.gov/pubmed/8976213>
319. Van Savage, J.G., et al. Transverse retubularized sigmoidovesicostomy continent urinary diversion to the umbilicus. *J Urol*, 2001. 166: 644. <http://www.ncbi.nlm.nih.gov/pubmed/11458110>
320. Karsenty, G., et al. A novel technique to achieve cutaneous continent urinary diversion in spinal cord-injured patients unable to catheterize through native urethra. *Spinal Cord*, 2008. 46: 305. <http://www.ncbi.nlm.nih.gov/pubmed/17700513>
321. Peterson, A.C., et al. Urinary diversion in patients with spinal cord injury in the United States. *Urology*, 2012. 80: 1247. <http://www.ncbi.nlm.nih.gov/pubmed/23206770>
322. Vanni, A.J., et al. Ileovesicostomy for the neurogenic bladder patient: outcome and cost comparison of open and robotic assisted techniques. *Urology*, 2011. 77: 1375. <http://www.ncbi.nlm.nih.gov/pubmed/21146864>
323. Wiener, J.S., et al. Bladder augmentation versus urinary diversion in patients with spina bifida in the United States. *J Urol*, 2011. 186: 161. <http://www.ncbi.nlm.nih.gov/pubmed/21575969>
324. Atan, A., et al. Advantages and risks of ileovesicostomy for the management of neuropathic bladder. *Urology*, 1999. 54: 636. <http://www.ncbi.nlm.nih.gov/pubmed/10510920>
325. Cass, A.S., et al. A 22-year followup of ileal conduits in children with a neurogenic bladder. *J Urol*, 1984. 132: 529. <http://www.ncbi.nlm.nih.gov/pubmed/6471190>



326. Hald, T., et al. Vesicostomy-an alternative urine diversion operation. Long term results. *Scand J Urol Nephrol*, 1978. 12: 227. <http://www.ncbi.nlm.nih.gov/pubmed/725543>
327. Schwartz, S.L., et al. Incontinent ileo-vesicostomy urinary diversion in the treatment of lower urinary tract dysfunction. *J Urol*, 1994. 152: 99. <http://www.ncbi.nlm.nih.gov/pubmed/8201699>
328. Herschorn, S., et al. Urinary undiversion in adults with myelodysplasia: long-term followup. *J Urol*, 1994. 152: <http://www.ncbi.nlm.nih.gov/pubmed/8015064> 329. <http://www.ncbi.nlm.nih.gov/pubmed/8015064>
329. Vasudeva, P., et al. Factors implicated in pathogenesis of urinary tract infections in neurogenic bladders: some revered, few forgotten, others ignored. *Neurourol Urodyn*, 2014. 33: 95. <http://www.ncbi.nlm.nih.gov/pubmed/23460489>
330. Bakke, A., et al. Bacteriuria in patients treated with clean intermittent catheterization. *Scand J Infect Dis*, 1991. 23: 577. <http://www.ncbi.nlm.nih.gov/pubmed/1767253>
331. Waites, K.B., et al. Epidemiology and risk factors for urinary tract infection following spinal cord injury. *Arch Phys Med Rehabil*, 1993. 74:691. <http://www.ncbi.nlm.nih.gov/pubmed/8328888>
332. Nicolle, L.E., et al. Infectious Diseases Society of America guidelines for the diagnosis and treatment of asymptomatic bacteriuria in adults. *Clin Infect Dis*, 2005. 40: 643. <http://www.ncbi.nlm.nih.gov/pubmed/15714408>
333. Goetz, L.L., et al. International Spinal Cord Injury Urinary Tract Infection Basic Data Set. *Spinal Cord*, 2013. 51: 700. <http://www.ncbi.nlm.nih.gov/pubmed/23896666>
334. Pannek, J. Treatment of urinary tract infection in persons with spinal cord injury: guidelines, evidence, and clinical practice. A questionnaire-based survey and review of the literature. *J Spinal Cord Med*, 2011. 34:11. <http://www.ncbi.nlm.nih.gov/pubmed/21528621>
335. Deville, W.L., et al. The urine dipstick test useful to rule out infections. A meta-analysis of the accuracy. *BMC Urol*, 2004. 4: 4. <http://www.ncbi.nlm.nih.gov/pubmed/15175113>
336. Hoffman, J.M., et al. Nitrite and leukocyte dipstick testing for urinary tract infection in individuals with spinal cord injury. *J Spinal Cord Med*, 2004. 27: 128. <http://www.ncbi.nlm.nih.gov/pubmed/15162883>
337. Biering-Sorensen, F., et al. Urinary tract infections in patients with spinal cord lesions: treatment and prevention. *Drugs*, 2001. 61: 1275. <http://www.ncbi.nlm.nih.gov/pubmed/11511022>
338. Everaert, K., et al. Urinary tract infections in spinal cord injury: prevention and treatment guidelines. *Acta Clin Belg*, 2009. 64:335. <http://www.ncbi.nlm.nih.gov/pubmed/19810421>
339. D'Hondt, F., et al. Urinary tract infections in patients with spinal cord injuries. *Curr Infect Dis Rep*, 2011. 13: 544. <http://www.ncbi.nlm.nih.gov/pubmed/21853416>
340. Jia, C., et al. Detrusor botulinum toxin A injection significantly decreased urinary tract infection in patients with traumatic spinal cord injury. *Spinal Cord*, 2013. 51: 487. <http://www.ncbi.nlm.nih.gov/pubmed/23357928>
341. Li, L., et al. Impact of hydrophilic catheters on urinary tract infections in people with spinal cord injury: systematic review and meta-analysis of randomized controlled trials. *Arch Phys Med Rehabil*, 2013. 94: 782. <http://www.ncbi.nlm.nih.gov/pubmed/23168400>
342. Waites, K.B., et al. Evaluation of 3 methods of bladder irrigation to treat bacteriuria in persons with neurogenic bladder. *J Spinal Cord Med*, 2006. 29: 217. <http://www.ncbi.nlm.nih.gov/pubmed/16859225>
343. Gallien, P., et al. Cranberry versus placebo in the prevention of urinary infections in multiple sclerosis: a multicenter, randomized, placebo-controlled, double-blind trial. *Mult Scler*, 2014. 20: 1252. <http://www.ncbi.nlm.nih.gov/pubmed/24402038>
344. Lee, B.S., et al. Methenamine hippurate for preventing urinary tract infections. *Cochrane Database Syst Rev*, 2012. 10: CD003265. <http://www.ncbi.nlm.nih.gov/pubmed/23076896>
345. Günther, M., et al. Harnwegsinfektprophylaxe. Urinansäuerung mittels L-Methionin bei neurogener Blasenfunktionsstörung. *Urologe B*, 2002. 42: 218.
346. Hachen, H.J. Oral immunotherapy in paraplegic patients with chronic urinary tract infections: a double-blind, placebo-controlled trial. *J Urol*, 1990. 143: 759. <http://www.ncbi.nlm.nih.gov/pubmed/2179584>
347. Salomon, J., et al. Prevention of urinary tract infection in spinal cord-injured patients: safety and efficacy of a weekly oral cyclic antibiotic (WOCA) programme with a 2 year follow-up-an observational prospective study. *J Antimicrob Chemother*, 2006. 57: 784. <http://www.ncbi.nlm.nih.gov/pubmed/16473921>
348. Darouiche, R.O., et al. Multicenter randomized controlled trial of bacterial interference for prevention of urinary tract infection in patients with neurogenic bladder. *Urology*, 2011. 78: 341. <http://www.ncbi.nlm.nih.gov/pubmed/21683991>
349. Pannek, J., et al. Usefulness of classical homoeopathy for the prevention of urinary tract infections in patients with neurogenic bladder dysfunction: A case series. *Indian J Res Homoeopathy*, 2014. 8: 31.
350. Rees, P.M., et al. Sexual function in men and women with neurological disorders. *Lancet*, 2007. 369: 512. <http://www.ncbi.nlm.nih.gov/pubmed/17292771>
351. Jungwirth, A., et al., EAU Guidelines on Male Infertility. In: *EAU Guidelines*, edition presented at the annual EAU Congress Munich 2016. ISBN 978-90-79754-98-4 <http://uroweb.org/guideline/male-infertility/>
352. Hatzimouratidis K., H., et al., EAU Guidelines on Male Sexual Dysfunction. In: *EAU Guidelines*, edition presented at the annual EAU Congress Munich 2016. ISBN 978-90-79754-98-4. <http://www.ncbi.nlm.nih.gov/pubmed/20189712>
353. Foley, F.W., *Sexuality, in Multiple Sclerosis: A Guide for Families* K. RC., Editor. 2006, Demos Medical Publishing: New York, USA.
354. Annon, J.S., *PLISSIT Therapy in Handbook of Innovative Psychotherapies.*, R. Corsini, Editor. 1981, Wiley & Sons: New York, USA.



355. Lombardi, G., et al. Treatments for erectile dysfunction in spinal cord patients: alternatives to phosphodiesterase type 5 inhibitors? A review study. *Spinal Cord*, 2015. <http://www.ncbi.nlm.nih.gov/pubmed/26193811>
356. Chen, L., et al. Phosphodiesterase 5 Inhibitors for the Treatment of Erectile Dysfunction: A Trade-off Network Meta-analysis. *Eur Urol*, 2015. 68: 674. <http://www.ncbi.nlm.nih.gov/pubmed/25817916>
357. Lombardi, G., et al. Ten years of phosphodiesterase type 5 inhibitors in spinal cord injured patients. *J Sex Med*, 2009. 6: 1248. <http://www.ncbi.nlm.nih.gov/pubmed/19210710>
358. Lombardi, G., et al. Treating erectile dysfunction and central neurological diseases with oral phosphodiesterase type 5 inhibitors. Review of the literature. *J Sex Med*, 2012. 9: 970. <http://www.ncbi.nlm.nih.gov/pubmed/22304626>
359. Cardenas, D.D., et al. Two phase 3, multicenter, randomized, placebo-controlled clinical trials of fampridine-SR for treatment of spasticity in chronic spinal cord injury. *Spinal Cord*, 2014. 52: 70. <http://www.ncbi.nlm.nih.gov/pubmed/24216616>
360. Strebler, R.T., et al. Apomorphine sublingual as primary or secondary treatment for erectile dysfunction in patients with spinal cord injury. *BJU Int*, 2004. 93: 100. <http://www.ncbi.nlm.nih.gov/pubmed/14678378>
361. Pohanka, M., et al. The long-lasting improvement of sexual dysfunction in patients with advanced, fluctuating Parkinson's disease induced by pergolide: evidence from the results of an open, prospective, one-year trial. *Parkinsonism Relat Disord*, 2005. 11: 509. <http://www.ncbi.nlm.nih.gov/pubmed/15994112>
362. Chancellor, M.B., et al. Prospective comparison of topical minoxidil to vacuum constriction device and intracorporeal papaverine injection in treatment of erectile dysfunction due to spinal cord injury. *Urology*, 1994. 43: 365. <http://www.ncbi.nlm.nih.gov/pubmed/8134992>
363. Cookson, M.S., et al. Long-term results with vacuum constriction device. *J Urol*, 1993. 149: 290. <http://www.ncbi.nlm.nih.gov/pubmed/8426404>
364. Denil, J., et al. Vacuum erection device in spinal cord injured men: patient and partner satisfaction. *Arch Phys Med Rehabil*, 1996. 77: 750. <http://www.ncbi.nlm.nih.gov/pubmed/8702367>
365. Levine, L.A. External devices for treatment of erectile dysfunction. *Endocrine*, 2004. 23: 157. <http://www.ncbi.nlm.nih.gov/pubmed/15146095>
366. Levine, L.A., et al. Vacuum constriction and external erection devices in erectile dysfunction. *Urol Clin North Am*, 2001. 28: 335. <http://www.ncbi.nlm.nih.gov/pubmed/11402585>
367. Bella, A.J., et al. Intracavernous pharmacotherapy for erectile dysfunction. *Endocrine*, 2004. 23: 149. <http://www.ncbi.nlm.nih.gov/pubmed/15146094>
368. Bodner, D.R., et al. The application of intracavernous injection of vasoactive medications for erection in men with spinal cord injury. *J Urol*, 1987. 138: 310. <http://www.ncbi.nlm.nih.gov/pubmed/3599245>
369. Dinsmore, W.W., et al. Treating men with predominantly nonpsychogenic erectile dysfunction with intracavernosal vasoactive intestinal polypeptide and phentolamine mesylate in a novel auto-injector system: a multicentre double-blind placebo-controlled study. *BJU Int*, 1999. 83: 274. <http://www.ncbi.nlm.nih.gov/pubmed/10233493>
370. Hirsch, I.H., et al. Use of intracavernous injection of prostaglandin E1 for neuropathic erectile dysfunction. *Paraplegia*, 1994. 32: 661. <http://www.ncbi.nlm.nih.gov/pubmed/7831071>
371. Kapoor, V.K., et al. Intracavernous papaverine for impotence in spinal cord injured patients. *Paraplegia*, 1993. 31: 675. <http://www.ncbi.nlm.nih.gov/pubmed/8259331>
372. Vidal, J., et al. Intracavernous pharmacotherapy for management of erectile dysfunction in multiple sclerosis patients. *Rev Neurol*, 1995. 23: 269. <http://www.ncbi.nlm.nih.gov/pubmed/7497173>
373. Deforge, D., et al. Male erectile dysfunction following spinal cord injury: a systematic review. *Spinal Cord*, 2006. 44: 465. <http://www.ncbi.nlm.nih.gov/pubmed/16317419>
374. Bodner, D.R., et al. Intraurethral alprostadil for treatment of erectile dysfunction in patients with spinal cord injury. *Urology*, 1999. 53: 199. <http://www.ncbi.nlm.nih.gov/pubmed/9886612>
375. Gross, A.J., et al. Penile prostheses in paraplegic men. *Br J Urol*, 1996. 78: 262. <http://www.ncbi.nlm.nih.gov/pubmed/8813925>
376. Kimoto, Y., et al. Penile prostheses for the management of the neuropathic bladder and sexual dysfunction in spinal cord injury patients: long term follow up. *Paraplegia*, 1994. 32: 336. <http://www.ncbi.nlm.nih.gov/pubmed/8058351>
377. Zermann, D.H., et al. Penile prosthetic surgery in neurologically impaired patients: long-term followup. *J Urol*, 2006. 175: 1041. <http://www.ncbi.nlm.nih.gov/pubmed/16469612>
378. Fode, M., et al. Male sexual dysfunction and infertility associated with neurological disorders. *Asian J Androl*, 2012. 14: 61. <http://www.ncbi.nlm.nih.gov/pubmed/22138899>
379. Lim, T.C., et al. A simple technique to prevent retrograde ejaculation during assisted ejaculation. *Paraplegia*, 1994. 32: 142. <http://www.ncbi.nlm.nih.gov/pubmed/8008416>
380. Philippon, M., et al. Successful pregnancies and healthy live births using frozen-thawed sperm retrieved by a new modified Hotchkiss procedure in males with retrograde ejaculation: first case series. *Basic Clin Androl*, 2015. 25: 5. <http://www.ncbi.nlm.nih.gov/pubmed/26034605>
381. Arafa, M.M., et al. Prostatic massage: a simple method of semen retrieval in men with spinal cord injury. *Int J Androl*, 2007. 30: 170. <http://www.ncbi.nlm.nih.gov/pubmed/17298549>
382. Kolettis, P.N., et al. Fertility outcomes after electroejaculation in men with spinal cord injury. *Fertil Steril*, 2002. 78: 429. <http://www.ncbi.nlm.nih.gov/pubmed/12137889>
383. Chehensse, C., et al. The spinal control of ejaculation revisited: a systematic review and metaanalysis of anejaculation in spinal cord injured patients. *Hum Reprod Update*, 2013. 19: 507. <http://www.ncbi.nlm.nih.gov/pubmed/23820516>



384. Beretta, G., et al. Reproductive aspects in spinal cord injured males. *Paraplegia*, 1989. 27: 113. <http://www.ncbi.nlm.nih.gov/pubmed/2717193>
385. Brackett, N.L., et al. Application of 2 vibrators salvages ejaculatory failures to 1 vibrator during penile vibratory stimulation in men with spinal cord injuries. *J Urol*, 2007. 177: 660. <http://www.ncbi.nlm.nih.gov/pubmed/17222653>
386. Sonksen, J., et al. Ejaculation induced by penile vibratory stimulation in men with spinal cord injuries. The importance of the vibratory amplitude. *Paraplegia*, 1994. 32: 651. <http://www.ncbi.nlm.nih.gov/pubmed/7831070>
387. Claydon, V.E., et al. Cardiovascular responses to vibrostimulation for sperm retrieval in men with spinal cord injury. *J Spinal Cord Med*, 2006. 29: 207. <http://www.ncbi.nlm.nih.gov/pubmed/16859224>
388. Eklund, M.B., et al. Incidence of autonomic dysreflexia and silent autonomic dysreflexia in men with spinal cord injury undergoing sperm retrieval: implications for clinical practice. *J Spinal Cord Med*, 2008. 31:33. <http://www.ncbi.nlm.nih.gov/pubmed/18533409>
389. Soler, J.M., et al. Midodrine improves ejaculation in spinal cord injured men. *J Urol*, 2007. 178: 2082. <http://www.ncbi.nlm.nih.gov/pubmed/17869290>
390. Pecori, C., et al. Paternal therapy with disease modifying drugs in multiple sclerosis and pregnancy outcomes: a prospective observational multicentric study. *BMC Neurol*, 2014. 14: 114. <http://www.ncbi.nlm.nih.gov/pubmed/24884599>
391. Brackett, N.L., et al. Treatment of infertility in men with spinal cord injury. *Nat Rev Urol*, 2010. 7: 162. <http://www.ncbi.nlm.nih.gov/pubmed/20157304>
392. Raviv, G., et al. Testicular sperm retrieval and intra cytoplasmic sperm injection provide favorable outcome in spinal cord injury patients, failing conservative reproductive treatment. *Spinal Cord*, 2013. 51: 642. <http://www.ncbi.nlm.nih.gov/pubmed/23689394>
393. Schatte, E.C., et al. Treatment of infertility due to anejaculation in the male with electroejaculation and intracytoplasmic sperm injection. *J Urol*, 2000. 163: 1717. <http://www.ncbi.nlm.nih.gov/pubmed/10799167>
394. Shieh, J.Y., et al. A protocol of electroejaculation and systematic assisted reproductive technology achieved high efficiency and efficacy for pregnancy for anejaculatory men with spinal cord injury. *Arch Phys Med Rehabil*, 2003. 84: 535. <http://www.ncbi.nlm.nih.gov/pubmed/12690592>
395. Taylor, Z., et al. Contribution of the assisted reproductive technologies to fertility in males suffering spinal cord injury. *Aust N Z J Obstet Gynaecol*, 1999. 39: 84. <http://www.ncbi.nlm.nih.gov/pubmed/10099757>
396. Rutkowski, S.B., et al. The influence of bladder management on fertility in spinal cord injured males. *Paraplegia*, 1995. 33: 263. <http://www.ncbi.nlm.nih.gov/pubmed/7630651>
397. Hamed, S.A., et al. Seminal fluid analysis and testicular volume in adults with epilepsy receiving valproate. *J Clin Neurosci*, 2015. 22: 508. <http://www.ncbi.nlm.nih.gov/pubmed/25636832>
398. Ohl, D.A., et al. Electroejaculation versus vibratory stimulation in spinal cord injured men: sperm quality and patient preference. *J Urol*, 1997. 157: 2147. <http://www.ncbi.nlm.nih.gov/pubmed/9146603>
399. Brackett, N.L., et al. Semen quality of spinal cord injured men is better when obtained by vibratory stimulation versus electroejaculation. *J Urol*, 1997. 157: 151. <http://www.ncbi.nlm.nih.gov/pubmed/8976239>
400. Brackett, N.L., et al. Semen retrieval in men with spinal cord injury is improved by interrupting current delivery during electroejaculation. *J Urol*, 2002. 167: 201. <http://www.ncbi.nlm.nih.gov/pubmed/11743305>
401. DeForge, D., et al. Fertility following spinal cord injury: a systematic review. *Spinal Cord*, 2005. 43: 693. <http://www.ncbi.nlm.nih.gov/pubmed/15951744>
402. Ferreiro-Velasco, M.E., et al. Sexual issues in a sample of women with spinal cord injury. *Spinal Cord*, 2005. 43: 51. <http://www.ncbi.nlm.nih.gov/pubmed/15303115>
403. Kreuter, M., et al. Sexuality and sexual life in women with spinal cord injury: a controlled study. *J Rehabil Med*, 2008. 40: 61. <http://www.ncbi.nlm.nih.gov/pubmed/18176739>
404. Kreuter, M., et al. Sexual adjustment and quality of relationship in spinal paraplegia: a controlled study. *Arch Phys Med Rehabil*, 1996. 77: 541. <http://www.ncbi.nlm.nih.gov/pubmed/8831469>
405. Kessler, T.M., et al. Sexual dysfunction in multiple sclerosis. *Expert Rev Neurother*, 2009. 9: 341. <http://www.ncbi.nlm.nih.gov/pubmed/19271943>
406. Lew-Starowicz, M., et al. Prevalence of Sexual Dysfunctions Among Women with Multiple Sclerosis. *Sex Disabil*, 2013. 31: 141. <http://www.ncbi.nlm.nih.gov/pubmed/23704801>
407. Reitz, A., et al. Impact of spinal cord injury on sexual health and quality of life. *Int J Impot Res*, 2004. 16:167. <http://www.ncbi.nlm.nih.gov/pubmed/14973522>
408. Harrison, J., et al. Factors associated with sexual functioning in women following spinal cord injury. *Paraplegia*, 1995. 33: 687. <http://www.ncbi.nlm.nih.gov/pubmed/8927405>
409. Westgren, N., et al. Sexuality in women with traumatic spinal cord injury. *Acta Obstet Gynecol Scand*, 1997. 76: 977. <http://www.ncbi.nlm.nih.gov/pubmed/9435740>
410. Lombardi, G., et al. Management of sexual dysfunction due to central nervous system disorders: a systematic review. *BJU Int*, 2015. 115 Suppl 6: 47. <http://www.ncbi.nlm.nih.gov/pubmed/25599613>
411. Fruhauf, S., et al. Efficacy of psychological interventions for sexual dysfunction: a systematic review and meta-analysis. *Arch Sex Behav*, 2013. 42: 915. <http://www.ncbi.nlm.nih.gov/pubmed/23559141>
412. Alexander, M., et al. Spinal cord injuries and orgasm: a review. *J Sex Marital Ther*, 2008. 34: 308. <http://www.ncbi.nlm.nih.gov/pubmed/18576233>
413. Sipski, M.L., et al. Sexual arousal and orgasm in women: effects of spinal cord injury. *Ann Neurol*, 2001. 49: 35. <http://www.ncbi.nlm.nih.gov/pubmed/11198294>



414. Sipski, M.L., et al. Physiologic parameters associated with sexual arousal in women with incomplete spinal cord injuries. *Arch Phys Med Rehabil*, 1997. 78: 305. <http://www.ncbi.nlm.nih.gov/pubmed/9084355>
415. McAlonan, S. Improving sexual rehabilitation services: the patient's perspective. *Am J Occup Ther*, 1996. 50: 826. <http://www.ncbi.nlm.nih.gov/pubmed/8947375>
416. Schopp, L.H., et al. Impact of comprehensive gynecologic services on health maintenance behaviours among women with spinal cord injury. *Disabil Rehabil*, 2002. 24: 899. <http://www.ncbi.nlm.nih.gov/pubmed/12519485>
417. Sukumaran, S.C., et al. Polytherapy increases the risk of infertility in women with epilepsy. *Neurology*, 2010. 75: 1351. <http://www.ncbi.nlm.nih.gov/pubmed/20938026>
418. Axel, S.J. Spinal cord injured women's concerns: menstruation and pregnancy. *Rehabil Nurs*, 1982. 7: 10. <http://www.ncbi.nlm.nih.gov/pubmed/6921826>
419. Jackson, A.B., et al. A multicenter study of women's self-reported reproductive health after spinal cord injury. *Arch Phys Med Rehabil*, 1999. 80: 1420. <http://www.ncbi.nlm.nih.gov/pubmed/10569436>
420. Baker, E.R., et al. Pregnancy in spinal cord injured women. *Arch Phys Med Rehabil*, 1996. 77: 501. <http://www.ncbi.nlm.nih.gov/pubmed/8629929>
421. Baker, E.R., et al. Risks associated with pregnancy in spinal cord-injured women. *Obstet Gynecol*, 1992. 80: 425. <http://www.ncbi.nlm.nih.gov/pubmed/1495699>
422. Cross, L.L., et al. Pregnancy, labor and delivery post spinal cord injury. *Paraplegia*, 1992. 30: 890. <http://www.ncbi.nlm.nih.gov/pubmed/1287543>
423. Hughes, S.J., et al. Management of the pregnant woman with spinal cord injuries. *Br J Obstet Gynaecol*, 1991. 98: 513. <http://www.ncbi.nlm.nih.gov/pubmed/1873238>
424. Dannels, A., et al. The perimenopause experience for women with spinal cord injuries. *SCI Nurs*, 2004. 21: 9. <http://www.ncbi.nlm.nih.gov/pubmed/15176344>
425. Vukusic, S., et al. Multiple sclerosis and pregnancy in the 'treatment era'. *Nat Rev Neurol*, 2015. 11: 280. <http://www.ncbi.nlm.nih.gov/pubmed/25896084>
426. Bove, R., et al. Management of multiple sclerosis during pregnancy and the reproductive years: a systematic review. *Obstet Gynecol*, 2014. 124: 1157. <http://www.ncbi.nlm.nih.gov/pubmed/25415167>
427. Austin, P.F., et al. The standardization of terminology of lower urinary tract function in children and adolescents: Update report from the standardization committee of the International Children's Continence Society. *J Urol*, 2014. PAEDIATRIC UROLOGY - UPDATE MARCH 2016 107. <http://www.ncbi.nlm.nih.gov/pubmed/24508614>
428. Hellstrom, A.L., et al. Micturition habits and incontinence in 7-year-old Swedish school entrants. *Eur J Pediatr*, 1990. 149: 434. <http://www.ncbi.nlm.nih.gov/pubmed/2332015>
429. Bakker, E., et al. Voiding habits and wetting in a population of 4,332 Belgian schoolchildren aged between 10 and 14 years. *Scand J Urol Nephrol*, 2002. 36: 354. <http://www.ncbi.nlm.nih.gov/pubmed/12487740>
430. Söderstrom, U., et al. Urinary and faecal incontinence: a population-based study. *Acta Paediatr*, 2004. 93: 386. <http://www.ncbi.nlm.nih.gov/pubmed/15124844>
431. Sureshkumar, P., et al. Daytime urinary incontinence in primary school children: a population-based survey. *J Pediatr*, 2000. 137: 814. <http://www.ncbi.nlm.nih.gov/pubmed/19110268>
432. Sureshkumar, P., et al. A population based study of 2,856 school-age children with urinary incontinence. *J Urol*, 2009. 181: 808. <http://www.ncbi.nlm.nih.gov/pubmed/11113838>
433. Veiga, M.L., et al. Constipation in children with isolated overactive bladders. *J Pediatr Urol*, 2013. 9: 945. <http://www.ncbi.nlm.nih.gov/pubmed/23462384>
434. Borch, L., et al. Bladder and bowel dysfunction and the resolution of urinary incontinence with successful management of bowel symptoms in children. *Acta Paediatr*, 2013. 102: e215. <http://www.ncbi.nlm.nih.gov/pubmed/23368903>
435. Chang, S.J., et al. Treatment of daytime urinary incontinence: A standardization document from the International Children's Continence Society. *Neurourol Urodyn*, 2015. <http://www.ncbi.nlm.nih.gov/pubmed/26473630>
436. Hoebeker, P., et al. Diagnostic evaluation of children with daytime incontinence. *J Urol*, 2010. 183: 699. <http://www.ncbi.nlm.nih.gov/pubmed/20022025>
437. Akbal, C., et al. Dysfunctional voiding and incontinence scoring system: quantitative evaluation of incontinence symptoms in pediatric population. *J Urol*, 2005. 173: 969. <http://www.ncbi.nlm.nih.gov/pubmed/15711352>
438. Farhat, W., et al. The dysfunctional voiding scoring system: quantitative standardization of dysfunctional voiding symptoms in children. *J Urol*, 2000. 164: 1011. <http://www.ncbi.nlm.nih.gov/pubmed/10958730>
439. Chang, S.J., et al. Constipation is associated with incomplete bladder emptying in healthy children. *Neurourol Urodyn*, 2012. 31: 105. <http://www.ncbi.nlm.nih.gov/pubmed/22038844>
440. Burgers, R.E., et al. Management of functional constipation in children with lower urinary tract symptoms: report from the Standardization Committee of the International Children's Continence Society. *J Urol*, 2013. 190: 29. <http://www.ncbi.nlm.nih.gov/pubmed/23313210>
441. van Gool, J.D., et al. Multi-center randomized controlled trial of cognitive treatment, placebo, oxybutynin, bladder training, and pelvic floor training in children with functional urinary incontinence. *Neurourol Urodyn*, 2014. 33: 482. <http://www.ncbi.nlm.nih.gov/pubmed/23775924>
442. Hellstrom, A.L. Urotherapy in children with dysfunctional bladder. *Scand J Urol Nephrol Suppl*, 1992. 141: 106. <http://www.ncbi.nlm.nih.gov/pubmed/1609245>
443. Barroso, U., Jr., et al. Electrical stimulation for lower urinary tract dysfunction in children: a systematic review of the literature. *Neurourol Urodyn*, 2011. 30: 1429. <http://www.ncbi.nlm.nih.gov/pubmed/21717502>



445. Bower, W.F., et al. A review of non-invasive electro neuromodulation as an intervention for nonneurogenic bladder dysfunction in children. *Neurourol Urodyn*, 2004. 23: 63. <http://www.ncbi.nlm.nih.gov/pubmed/14694460>
446. De Paepe, H., et al. Pelvic-floor therapy in girls with recurrent urinary tract infections and dysfunctional voiding. *Br J Urol*, 1998. 81 Suppl 3: 109. <http://www.ncbi.nlm.nih.gov/pubmed/9634033>
447. Vijverberg, M.A., et al. Bladder rehabilitation, the effect of a cognitive training programme on urge incontinence. *Eur Urol*, 1997. 31: 68. 108 PAEDIATRIC UROLOGY - UPDATE MARCH 2016. <http://www.ncbi.nlm.nih.gov/pubmed/9032538>
448. Lordêlo, P., et al. Prospective study of transcutaneous parasacral electrical stimulation for overactive bladder in children: long-term results. *J Urol*, 2009. 182: 2900. <http://www.ncbi.nlm.nih.gov/pubmed/19846164>
449. Ladi-Seyedian, S., et al. Management of non-neuropathic underactive bladder in children with voiding dysfunction by animated biofeedback: a randomized clinical trial. *Urology*, 2015. 85: 205. <http://www.ncbi.nlm.nih.gov/pubmed/25444633>
450. Kajbafzadeh, A.M., et al. Transcutaneous interferential electrical stimulation for the management of non-neuropathic underactive bladder in children: a randomised clinical trial. *BJU Int*, 2015. <http://www.ncbi.nlm.nih.gov/pubmed/22065105>
451. Featherstone, N., et al. Ephedrine hydrochloride: novel use in the management of resistant nonneurogenic daytime urinary incontinence in children. *J Pediatr Urol*, 2013. 9: 915. <http://www.ncbi.nlm.nih.gov/pubmed/23332206>
452. Nijman, R.J., et al. Tolterodine treatment for children with symptoms of urinary urge incontinence suggestive of detrusor overactivity: results from 2 randomized, placebo controlled trials. *J Urol*, 2005. 173: 1334. <http://www.ncbi.nlm.nih.gov/pubmed/15758796>
453. Marschall-Kehrel, D., et al. Treatment with propiverine in children suffering from nonneurogenic overactive bladder and urinary incontinence: results of a randomized placebo-controlled phase 3 clinical trial. *Eur Urol*, 2009. 55: 729. <http://www.ncbi.nlm.nih.gov/pubmed/18502028>
454. Kramer, S.A., et al. Double-blind placebo controlled study of alpha-adrenergic receptor antagonists (doxazosin) for treatment of voiding dysfunction in the pediatric population. *J Urol*, 2005. 173: 2121. <http://www.ncbi.nlm.nih.gov/pubmed/15879863>
455. Hoebeke, P., et al. The effect of botulinum-A toxin in incontinent children with therapy resistant overactive detrusor. *J Urol*, 2006. 176: 328. <http://www.ncbi.nlm.nih.gov/pubmed/16753434>
456. Groen, L.A., et al. Sacral neuromodulation with an implantable pulse generator in children with lower urinary tract symptoms: 15-year experience. *J Urol*, 2012. 188: 1313. <http://www.ncbi.nlm.nih.gov/pubmed/22902022>